

10/511008

DT05 d PCT/PTO 12 OCT 2004

S & F Ref: 413151

AUSTRALIA

PATENTS ACT 1990

COMPLETE SPECIFICATION

FOR A STANDARD PATENT

ORIGINAL

Name and Address
of Applicant:

Macquarie Veterinary Supplies Pty Ltd
4 Charlotte Street
Red Hill A.C.T 2603
AUSTRALIA

Actual Inventor(s): John Russell Beckett

Address for Service: Spruson & Ferguson, Patent Attorneys
Level 33 St Martins Tower, 31 Market Street
Sydney, New South Wales, 2000, Australia

Invention Title: An Aqueous Metal Bicarbonate Solution and Method of Use

ASSOCIATED PROVISIONAL APPLICATION DETAILS

[31] Application No(s)
P05677
PP0608

[33] Country
AU
AU

[32] Application Date
17 March 1997
28 November 1997

The following statement is a full description of this invention,
including the best method of performing it known to me/us:-

An Aqueous Metal Bicarbonate Solution and Method of Use

Technical Field

This invention relates to an aqueous metal bicarbonate solution, a process of preparing the aqueous metal bicarbonate solution and a method of preventing and treating certain
5 inflammatory diseases, degenerative diseases and viral diseases in mammals.

Generally the certain inflammatory diseases, degenerative diseases and viral diseases in mammals are those that require extracellular or intracellular acidic conditions or extracellular or intracellular proton concentrations at some point in disease process or disease pathogenesis.

10 Typically the certain inflammatory diseases, degenerative diseases and viral diseases in mammals are those that require the activities of carbonic anhydrase enzymes and/or the activities of acid (aspartic) protease enzymes and/or the activities of endosomal or lysosomal acid-requiring-enzymes and/or the activities of V-type ATPase proton pumps at
15 some point in disease process or disease pathogenesis. Typically the certain inflammatory diseases, degenerative diseases and viral diseases in mammals are represented by the diseases of arthritis and influenza.

This invention relates to a method of using an aqueous metal bicarbonate solution to decrease senescence and to increase longevity in mammals. Generally senescence is decreased and longevity is increased in mammals by improving the buffering capacity of
20 the extracellular and intracellular fluids of the body. Generally senescence is decreased and longevity is increased in mammals by the improved buffering capacity causing a decrease in proton concentrations in the extracellular and intracellular fluids of the body. Typically senescence is decreased and longevity is increased in mammals by improving the buffering capacity of the extracellular and intracellular bicarbonate buffers. Typically
25 senescence is decreased and longevity is increased in mammals by the improved extracellular and intracellular bicarbonate buffers causing a decrease in proton concentrations.

Background Art

Certain inflammatory diseases, degenerative diseases and viral diseases are major causes
30 of morbidity and mortality in mammals. Typically these diseases are represented by the diseases of arthritis and influenza.

Arthritis is any inflammatory condition of the joints, characterised by pain and swelling. Osteoarthritis is the most common form of arthritis in which one or many joints undergo degenerative changes. Treatment includes rest of the involved joints, heat, and antiinflammatory drugs. Intraarticular injections of corticosteroids may give relief.

5 Surgical treatment is sometimes necessary and may reduce pain and greatly improve the function of the joint. However these treatments, apart from surgical treatment, only provide temporary relief and some may have severe side reactions.

Influenza is a highly contagious infection of the respiratory tract caused by a myxovirus and transmitted by airborne droplet infection. It occurs in isolated cases, epidemics and
10 pandemics. Treatment is symptomatic and usually involves bed rest, antipyretics such as aspirin and drinking of fluids. New strains of the virus emerge at regular intervals so it is difficult to take preventative measures to avoid the infection. There is a need for a method to prevent and to treat certain inflammatory diseases, degenerative diseases and viral diseases in mammals. There is a need for a method to prevent and to treat arthritis and
15 influenza in mammals.

Senescence in mammals is characterised by progressive oxidations of the structural and functional molecules that constitute body cells and tissues. Oxidations of the structural and functional molecules in body cells and tissues are increased in rate by acidic conditions. Oxidations of structural and functional molecules are increased in rate by the presence of
20 excess proton concentrations. There is a need for a method to prevent and treat excess proton concentrations in body cells so that oxidations of structural and functional molecules are decreased in rate. There is a need for a method to decrease and treat senescence in mammals.

Objects of Invention

25 It is an object of this invention to provide an aqueous metal bicarbonate solution to prevent and to treat certain inflammatory diseases, degenerative diseases and viral diseases in mammals. It is a further object of this invention to provide a process of preparing the aqueous metal bicarbonate solution. It is also an object of this invention to provide methods for the prevention and treatment of certain inflammatory diseases, degenerative
30 diseases and viral diseases in mammals using the aqueous metal bicarbonate solution. Generally the certain inflammatory diseases, degenerative diseases and viral diseases in mammals are those that require extracellular or intracellular acidic conditions or

extracellular or intracellular proton concentrations at some point in disease process or disease pathogenesis. Typically the certain inflammatory diseases, degenerative diseases and viral diseases in mammals are those that require the activities of carbonic anhydrase enzymes and/or the activities of acid (aspartic) protease enzymes and/or the activities of endosomal or lysosomal acid-requiring-enzymes and/or the activities of V-type ATPase proton pumps at some point in disease process or disease pathogenesis. Typically the certain inflammatory diseases, degenerative diseases and viral diseases in mammals are represented by the diseases of arthritis and influenza.

It is an object of this invention to provide an aqueous metal bicarbonate solution to decrease senescence and to treat senescence and to increase longevity in mammals. It is a further object of this invention to provide a process of preparing the aqueous metal bicarbonate solution. It is also an object of this invention to provide methods for the decrease of senescence and the treatment of senescence and the increase in longevity in mammals using the aqueous metal bicarbonate solution. Generally senescence is decreased and longevity is increased in mammals by improving the buffering capacity of the extracellular and intracellular fluids of the body. Generally senescence is decreased and longevity is increased in mammals by the improved buffering capacity causing a decrease in proton concentrations in the extracellular and intracellular fluids of the body. Typically senescence is decreased and longevity is increased in mammals by improving the buffering capacity of the extracellular and intracellular bicarbonate buffers. Typically senescence is decreased and longevity is increased in mammals by the improved extracellular and intracellular bicarbonate buffers causing a decrease in proton concentrations.

The aim of the invention is to achieve at least one of the stated objects.

Disclosure of Invention

The specification describes an aqueous metal bicarbonate solution comprising a stoichiometric concentration of bicarbonate anions and a corresponding substantially stoichiometric concentration of metal cations in association with the bicarbonate anions, the metal bicarbonate being present in a therapeutically effective amount and an acceptable carbon dioxide-containing-aqueous diluent to maintain the metal bicarbonate in the aqueous diluent.

Typically the solution is acceptable for oral administration.

In one embodiment there is provided a combination comprising the described solution in combination with a stabilising agent in an amount effective to maintain and



stabilise the bicarbonate anions in the neutral to mildly alkaline solution. Generally the combination is kept in a sealed or closed container at 0.8 to 5 atmospheres, more typically 1 atmosphere at 0 - 25°C, more typically 0. 1-10°C. In one particular embodiment the stabilising agent may be present in the solution in an amount effective to maintain and

5 stabilise the bicarbonate anions in the neutral to mildly alkaline solution. In another particular embodiment the stabilising agent may consist of or comprise a gas above the solution in an amount effective to maintain and stabilise the bicarbonate anions in the neutral to mildly alkaline solution. In a further particular embodiment the stabilising agent may be present in the solution and also may consist of or comprise a gas above the

10 solution, the total amount of stabilising agent in the solution and in the gas above the solution being in an amount effective to maintain and stabilise the bicarbonate anions in the neutral to mildly alkaline solution. The stabilising agent which consists of a gas above the solution may be carbon dioxide. The stabilising agent which comprises a gas above the solution may be carbon dioxide in an inert gas such as nitrogen, air, oxygen,

15 argon and/or helium, for example. The stabilising agent in the solution may be carbon dioxide dissolved in the solution, hydrated carbon dioxide, carbonic acid, and/or other suitable source of carbon dioxide.

The specification describes a process of preparing an aqueous metal bicarbonate solution comprising a stoichiometric concentration of bicarbonate anions and a

20 corresponding substantially stoichiometric concentration of metal cations in association with the bicarbonate anions, which process comprises reacting a concentration of a metal carbonate or metal carbonate hydroxide or metal oxide or metal hydroxide with a concentration of carbonic acid or hydrated carbon dioxide to produce the metal bicarbonate aqueous solution, wherein said metal bicarbonate being present in a

25 therapeutically effective amount.

Typically the aqueous metal bicarbonate solution has a neutral to mildly alkaline pH. Typically the pH is in the range 7 to 9. Typically the temperature of the aqueous metal bicarbonate solution is maintained at a level to maintain the metal bicarbonate in the aqueous diluent.

30 The specification describes an aqueous metal bicarbonate solution whenever prepared by the described process.

The specification describes a method of preventing and treating certain inflammatory diseases and degenerative diseases in a mammal in need of such prevention or treatment comprising administering to said mammal an effective amount of the

35 described aqueous metal bicarbonate solution or a metal bicarbonate.



Generally the certain inflammatory diseases and degenerative diseases in a mammal are those that require extracellular or intracellular acidic conditions or extracellular or intracellular proton concentrations at some point in disease process or disease pathogenesis.

5 Typically the certain inflammatory diseases and degenerative diseases in a mammal are those that require the activities of carbonic anhydrase enzymes and/or the activities of acid (aspartic) protease enzymes and/or the activities of endosomal or lysosomal acid-requiring-enzymes and/or the activities of V-type ATPase proton pumps at some point in disease process or disease pathogenesis.

10 Typically the certain inflammatory diseases or degenerative diseases may present as arthritis. Typically the arthritis may present as osteoarthritis.

The specification describes a method of preventing and treating certain viral diseases in a mammal in need of such prevention or treatment comprising administering to said mammal an effective amount of the described aqueous metal bicarbonate solution or a metal bicarbonate.

15 Typically the certain viral diseases require intracellular acidic conditions or intracellular proton concentrations for either removal of viral protein coats or assembly of viral protein coats. Typically the viral diseases may present as influenza.

The specification describes a method of decreasing and treating senescence and of increasing longevity in a mammal comprising administering to said mammal an effective amount of the described aqueous metal bicarbonate solution or a metal bicarbonate.

20 Typically senescence is decreased and longevity is increased by maintaining or increasing normal extracellular and/or intracellular alkaline conditions. Typically senescence is decreased and longevity is increased by improving the buffering capacity of the extracellular and intracellular fluids of the body. Typically longevity is increased by maintaining or increasing normal mitochondrial alkaline conditions. Typically longevity is increased by decreasing extracellular and intracellular acidic conditions or by decreasing extracellular and intracellular proton concentrations.

25 Typically senescence is decreased and longevity is increased in mammals by improving the buffering capacity of the extracellular and intracellular bicarbonate buffers. Typically senescence is decreased and longevity is increased in mammals by the improved extracellular and intracellular bicarbonate buffers causing a decrease in proton concentrations. Typically senescence is decreased and longevity is increased by preventing or treating certain inflammatory diseases, degenerative diseases and viral



diseases in mammals. Typically longevity is increased by decreasing the morbidity and mortality associated with these diseases.

The specification describes a method of scavenging protons in a mammal comprising administering to said mammal an effective amount of a proton scavenger.

5 Typically the proton scavenger comprises a metal bicarbonate. Typically the metal bicarbonate is in the form of the described aqueous metal bicarbonate solution.

The specification describes a method of decreasing proton concentrations in a mammal by altering carbonic anhydrase enzyme reactions in said mammal comprising administering to said mammal an effective amount of the described aqueous metal
10 bicarbonate solution or a metal bicarbonate.

The specification describes a method of decreasing inflammation and inflammatory conditions in a mammal comprising administering to said mammal an effective amount of the described aqueous metal bicarbonate solution or a metal bicarbonate.

15 Generally inflammation and inflammatory conditions are decreased by decreasing the extracellular and intracellular acidic conditions that are required for inflammatory processes. Generally inflammation and inflammatory conditions are decreased by decreasing the extracellular and intracellular proton concentrations that are required for inflammatory processes. Typically inflammation is decreased by altering carbonic
20 anhydrase enzyme reactions and/or decreasing the activities of acid (aspartic) protease enzymes and/or decreasing the activities of endosomal or lysosomal acid-requiring-enzymes and/or decreasing the activities of V-type ATPase proton pumps.

The specification describes a method of increasing motor activity in a mammal comprising administering to said mammal an effective amount of the described aqueous metal bicarbonate solution or a metal bicarbonate.

25 Typically motor activity is increased by decreasing extracellular and intracellular acidic conditions or by decreasing extracellular and intracellular proton concentrations. Typically motor activity is increased by improving the buffering capacity of extracellular and intracellular fluids. Typically motor activity is increased by improving the buffering capacity of the extracellular and intracellular bicarbonate buffers. Typically motor
30 activity is increased by increasing extracellular and intracellular alkaline conditions. Typically motor activity is increased by scavenging protons produced by ATP hydrolysis, lactic acid production, lipid metabolism and other metabolic processes.

According to a first embodiment of the present invention there is provided an aqueous neutral to mildly alkaline metal bicarbonate solution, comprising magnesium bicarbonate dissolved in the solution, said magnesium bicarbonate comprising



bicarbonate anions and magnesium cations, and a pH adjusting agent in the solution in an amount whereby the solution is at a neutral to mildly alkaline pH.

Typically a corresponding substantially stoichiometric concentration of metal cations are in association with the bicarbonate anions. Typically the solution is acceptable for oral administration.

In one embodiment there is provided a combination comprising a substantially stable aqueous neutral to mildly alkaline metal bicarbonate solution, comprising metal bicarbonate dissolved in the solution, said metal bicarbonate comprising bicarbonate anions and metal cations, and a pH adjusting agent in the solution in an amount whereby the solution is at a neutral to mildly alkaline pH, in combination with a stabilising agent in an amount effective to maintain and stabilise the bicarbonate anions in the neutral to mildly alkaline solution. In another embodiment there is provided a combination comprising a substantially stable aqueous neutral to mildly alkaline metal bicarbonate solution, comprising metal bicarbonate dissolved in the solution, said metal bicarbonate comprising bicarbonate anions and metal cations, in combination with a stabilising agent in an amount effective to maintain and stabilise the bicarbonate anions in the solution whereby the solution is at a neutral to mildly alkaline pH.

The pH adjusting agent and the stabilising agent may be the same or different. Generally the combination is kept in a sealed or closed container at 0.8 to 5 atmospheres, more typically 1 atmosphere at 0 - 25°C, more typically 0.1-10°C.

In one particular embodiment the stabilising agent may be present in the solution in an amount effective to maintain and stabilise the bicarbonate anions in the neutral to mildly alkaline solution. In another particular embodiment the stabilising agent may consist of or comprise a gas above the solution in an amount effective to maintain and stabilise the bicarbonate anions in the neutral to mildly alkaline solution. In a further particular embodiment the stabilising agent may be present in the solution and may consist of or comprise a gas above the solution, the total amount of stabilising agent in the solution and in the gas above the solution being in an amount effective to maintain and stabilise the bicarbonate anions in the neutral to mildly alkaline solution. The stabilising agent which consists of a gas above the solution may be carbon dioxide. The stabilising agent which comprises a gas above the solution may be carbon dioxide in an inert gas such as nitrogen, air, oxygen, argon and/or helium, for example. The stabilising agent in the solution may be carbon dioxide dissolved in the solution, hydrated carbon dioxide, carbonic acid, and/or other suitable source of carbon dioxide.



According to a second embodiment of the present invention there is provided a solution for preventing certain inflammatory diseases in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of the first embodiment wherein the magnesium bicarbonate is in an amount effective to prevent said diseases.

5 According to a third embodiment of the present invention there is provided a solution for treating certain inflammatory diseases in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of the first embodiment wherein the magnesium bicarbonate is in an amount effective to treat said diseases.

10 According to a fourth embodiment of the present invention there is provided a solution for preventing degenerative diseases in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of the first embodiment wherein the magnesium bicarbonate is in an amount effective to prevent said diseases.

15 According to a fifth embodiment of the present invention there is provided a solution for treating degenerative diseases in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of the first embodiment wherein the magnesium bicarbonate is in an amount effective to treat said diseases.

20 According to a sixth embodiment of the present invention there is provided a solution for preventing certain viral diseases in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of the first embodiment wherein the magnesium bicarbonate is in an amount effective to prevent said diseases.

According to a seventh embodiment of the present invention there is provided a solution for treating certain viral diseases in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of the first embodiment wherein the magnesium bicarbonate is in an amount effective to treat said diseases.

25 According to an eighth embodiment of the present invention there is provided a solution for decreasing senescence in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of the first embodiment wherein the magnesium bicarbonate is in an amount effective to decrease senescence.

30 According to a ninth embodiment of the present invention there is provided a solution for treating senescence in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of the first embodiment wherein the magnesium bicarbonate is in an amount effective to treat senescence.

According to a tenth embodiment of the present invention there is provided a solution for increasing longevity in a mammal, comprising the aqueous neutral to mildly



alkaline metal bicarbonate solution of the first embodiment wherein the magnesium bicarbonate is in an amount effective to increase longevity.

According to an eleventh embodiment of the present invention there is provided a solution for scavenging protons in a mammal, comprising the aqueous neutral to mildly
5 alkaline metal bicarbonate solution of the first embodiment wherein the magnesium bicarbonate is in an amount effective to scavenge protons.

According to a twelfth embodiment of the present invention there is provided a solution for decreasing proton concentrations in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of the first embodiment wherein the
10 magnesium bicarbonate is in an amount effective to decrease proton concentrations.

According to a thirteenth embodiment of the present invention there is provided a solution for decreasing inflammation in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of the first embodiment, wherein the magnesium bicarbonate is in an amount effective to decrease inflammation.

According to a fourteenth embodiment of the present invention there is provided a solution for decreasing inflammatory conditions in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of the first embodiment, wherein the magnesium bicarbonate is in an amount effective to decrease inflammatory conditions.
15

According to a fifteenth embodiment of the present invention there is provided a solution for increasing motor activity in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of the first embodiment wherein the magnesium bicarbonate is in an amount effective to increase motor activity.
20

According to a sixteenth embodiment of the present invention there is provided a solution for decreasing fatigue in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of the first embodiment wherein the magnesium bicarbonate is in an amount effective to decrease fatigue.
25

According to a seventeenth embodiment of the present invention there is provided a process of preparing an aqueous neutral to mildly alkaline metal bicarbonate solution comprising bicarbonate anions and metal cations, which process comprises reacting a compound selected from the group consisting of metal carbonate, metal carbonate hydroxide, metal oxide, metal hydroxide and any mixture thereof with an effective concentration of a pH adjusting agent to produce the aqueous neutral to mildly alkaline metal bicarbonate solution, wherein the pH adjusting agent is present in an amount
30 whereby the solution is at a neutral to mildly alkaline pH.



Typically a corresponding substantially stoichiometric concentration of metal cations are in association with the bicarbonate anions. Generally the solution is stored in a sealed or closed container at 0.8 to 5 atmospheres, more typically 1 atmosphere at 0 - 25°C, more typically 0.1-10°C. In one embodiment the process further comprises combining the solution with a stabilising agent in an amount effective to maintain and stabilise the bicarbonate anions in the neutral to mildly alkaline solution. In one particular embodiment the process comprises conducting the process under gaseous atmosphere comprising a stabilising agent in an amount effective to maintain and stabilise the bicarbonate anions in the neutral to mildly alkaline solution. The stabilising agent may be carbon dioxide or comprise carbon dioxide in an inert gas such as nitrogen, air, oxygen, argon and/or helium, for example. Generally the combination is stored in a sealed or closed container at 0.8 to 5 atmospheres, more typically 1 atmosphere at 0 - 25°C, more typically 0.1-10°C.

One particular embodiment may comprise adding the stabilising agent to the solution in the solution in an amount effective to maintain and stabilise the bicarbonate anions in the neutral to mildly alkaline solution. Another particular embodiment may comprise blanketing the solution with a gas consisting of or comprising the stabilising agent in an amount effective to maintain and stabilise the bicarbonate anions in the neutral to mildly alkaline solution. A further particular embodiment may comprise adding the stabilising agent to the solution in the solution and blanketing the solution with a gas consisting of or comprising the stabilising agent, the total amount of stabilising agent in the solution and in the gas above the solution being in an amount effective to maintain and stabilise the bicarbonate anions in the neutral to mildly alkaline solution. The stabilising agent which consists of a gas above the solution may be carbon dioxide. The stabilising agent which comprises a gas above the solution may be carbon dioxide in an inert gas such as nitrogen, air, oxygen, argon and/or helium, for example. The stabilising agent in the solution may be carbon dioxide dissolved in the solution, hydrated carbon dioxide, carbonic acid, and/or other suitable source of carbon dioxide.

According to an eighteenth embodiment of the present invention there is provided a aqueous neutral to mildly alkaline metal bicarbonate solution whenever prepared by the process of the seventeenth embodiment.

According to a nineteenth embodiment of the present invention there is provided a combination comprising a substantially stable aqueous neutral to mildly alkaline metal bicarbonate solution, comprising magnesium bicarbonate dissolved in the solution, said magnesium bicarbonate comprising bicarbonate anions and magnesium cations, and a pH



adjusting agent in the solution in an amount whereby the solution is at a neutral to mildly alkaline pH, in combination with a stabilising agent in an amount effective to maintain and stabilise the bicarbonate anions in the neutral to mildly alkaline solution.

According to a twentieth embodiment of the present invention there is provided a method of preventing certain inflammatory diseases in a mammal in need of such prevention comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the second embodiment or a combination of the nineteenth embodiment.

According to a twenty-first embodiment of the present invention there is provided a method of treating certain inflammatory diseases in a mammal in need of such treatment comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the third embodiment or a combination of the nineteenth embodiment.

According to a twenty-second embodiment of the present invention there is provided a method of preventing degenerative diseases in a mammal in need of such prevention comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the fourth embodiment or a combination of the nineteenth embodiment.

According to a twenty-third embodiment of the present invention there is provided a method of treating degenerative diseases in a mammal in need of such treatment comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the fifth embodiment or a combination of the nineteenth embodiment.

Generally the certain inflammatory diseases and degenerative diseases in a mammal are those that require extracellular or intracellular acidic conditions or extracellular or intracellular proton concentrations at some point in disease process or disease pathogenesis.

Typically the certain inflammatory diseases and degenerative diseases in a mammal are those that require the activities of carbonic anhydrase enzymes and/or the activities of acid (aspartic) protease enzymes and/or the activities of endosomal or lysosomal acid-requiring-enzymes and/or the activities of V-type ATPase proton pumps at some point in disease process or disease pathogenesis.

Typically the certain inflammatory diseases or degenerative diseases may present as arthritis. Typically the arthritis may present as osteoarthritis.



According to a twenty-fourth embodiment of the present invention there is provided a method of preventing certain viral diseases in a mammal in need of such prevention comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the sixth embodiment or a combination of the nineteenth embodiment.

According to a twenty-fifth embodiment of the present invention there is provided a method of treating certain viral diseases in a mammal in need of such treatment comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the seventh embodiment or a combination of the nineteenth embodiment.

Typically the certain viral diseases require intracellular acidic conditions or intracellular proton concentrations for either removal of viral protein coats or assembly of viral protein coats. Typically the viral diseases may present as influenza.

According to a twenty-sixth embodiment of the present invention there is provided a method of decreasing senescence in a mammal comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the eighth embodiment or a combination of the nineteenth embodiment.

According to an twenty-seventh embodiment of the present invention there is provided a method of treating senescence in a mammal comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the ninth embodiment or a combination of the nineteenth embodiment.

According to a twenty-eighth embodiment of the present invention there is provided a method of increasing longevity in a mammal comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the tenth embodiment or a combination of the nineteenth embodiment.

Typically senescence is decreased and longevity is increased by maintaining or increasing normal extracellular and/or intracellular alkaline conditions. Typically senescence is decreased and longevity is increased by improving the buffering capacity of the extracellular and intracellular fluids of the body. Typically longevity is increased by maintaining or increasing normal mitochondrial alkaline conditions. Typically longevity is increased by decreasing extracellular and intracellular acidic conditions or by decreasing extracellular and intracellular proton concentrations. Typically senescence is



decreased and longevity is increased in mammals by improving the buffering capacity of the extracellular and intracellular bicarbonate buffers. Typically senescence is decreased and longevity is increased in mammals by the improved extracellular and intracellular bicarbonate buffers causing a decrease in proton concentrations.

5 Typically senescence is decreased and longevity is increased by preventing or treating certain inflammatory diseases, degenerative diseases and viral diseases in mammals. Typically longevity is increased by decreasing the morbidity and mortality associated with these diseases.

10 According to a twenty-ninth embodiment of the present invention there is provided a method of scavenging protons in a mammal comprising administering to said mammal an effective amount of the aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the eleventh embodiment or a combination of the nineteenth embodiment.

15 According to a thirtieth embodiment of the present invention there is provided a method of decreasing proton concentrations in a mammal by altering carbonic anhydrase enzyme reactions in said mammal comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the twelfth embodiment or a combination of the nineteenth embodiment.

20 According to a thirty-first embodiment of the present invention there is provided a method of decreasing inflammation in a mammal comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the thirteenth embodiment or a combination of the nineteenth embodiment.

25 According to a thirty-second embodiment of the present invention there is provided a method of decreasing inflammatory conditions in a mammal comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the fourteenth embodiment or a combination of the nineteenth embodiment.

30 Generally inflammation and inflammatory conditions are decreased by decreasing the extracellular and intracellular acidic conditions that are required for inflammatory processes. Generally inflammation and inflammatory conditions are decreased by decreasing the extracellular and intracellular proton concentrations that are required for inflammatory processes. Typically inflammation is decreased by altering carbonic anhydrase enzyme reactions and/or decreasing the activities of acid (aspartic) protease



enzymes and/or decreasing the activities of endosomal or lysosomal acid-requiring-enzymes and/or decreasing the activities of V-type ATPase proton pumps.

According to a thirty-third embodiment of the present invention there is provided a method of increasing motor activity in a mammal comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the fifteenth embodiment or a combination of the nineteenth embodiment.

According to a thirty-fourth embodiment of the present invention there is provided a method of decreasing fatigue in a mammal comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the sixteenth embodiment or a combination of the nineteenth embodiment.

The methods of the invention typically involve orally administering to the mammal, the mammal being typically human. Further the methods of the invention typically involve orally administering to a mammal in need of treatment for the specified condition of the particular embodiment, the mammal being typically human. Typically motor activity is increased by decreasing extracellular and intracellular acidic conditions or by decreasing extracellular and intracellular proton concentrations. Typically motor activity is increased by improving the buffering capacity of extracellular and intracellular fluids. Typically motor activity is increased by improving the buffering capacity of the extracellular and intracellular bicarbonate buffers. Typically motor activity is increased by increasing extracellular and intracellular alkaline conditions. Typically motor activity is increased by scavenging protons produced by ATP hydrolysis, lactic acid production, lipid metabolism and other metabolic processes.

According to a thirty-fifth embodiment of the present invention there is provided use of bicarbonate, magnesium cations and a pH adjusting agent of the first or eighteenth embodiment or a combination of the nineteenth embodiment for the manufacture of a medicament to prevent certain inflammatory diseases in a mammal in need of such prevention.

According to a thirty-sixth embodiment of the present invention there is provided use of bicarbonate, magnesium cations and a pH adjusting agent of the first or eighteenth embodiment or a combination of the nineteenth embodiment for the manufacture of a medicament to treat certain inflammatory diseases in a mammal in need of such treatment.



According to a thirty-seventh embodiment of the present invention there is provided use of bicarbonate, magnesium cations and a pH adjusting agent of the first or eighteenth embodiment or a combination of the nineteenth embodiment for the manufacture of a medicament to prevent degenerative diseases in a mammal in need of such prevention.

5 According to a thirty-eighth embodiment of the present invention there is provided use of bicarbonate, magnesium cations and a pH adjusting agent of the first or eighteenth embodiment or a combination of the nineteenth embodiment for the manufacture of a medicament to treat degenerative diseases in a mammal in need of such treatment.

10 According to a thirty-ninth embodiment of the present invention there is provided use of bicarbonate, magnesium cations and a pH adjusting agent of the first or eighteenth embodiment or a combination of the nineteenth embodiment for the manufacture of a medicament to prevent certain viral diseases in a mammal in need of such prevention.

15 According to a fortieth embodiment of the present invention there is provided use of bicarbonate, magnesium cations and a pH adjusting agent of the first or eighteenth embodiment or a combination of the nineteenth embodiment for the manufacture of a medicament to treat certain viral diseases in a mammal in need of such treatment.

20 According to a forty-first embodiment of the present invention there is provided use of bicarbonate, magnesium cations and a pH adjusting agent of the first or eighteenth embodiment or a combination of the nineteenth embodiment for the manufacture of a medicament to decrease senescence in a mammal.

25 According to a forty-second embodiment of the present invention there is provided use of bicarbonate, magnesium cations and a pH adjusting agent of the first or eighteenth embodiment or a combination of the nineteenth embodiment for the manufacture of a medicament to treat senescence in a mammal.

30 According to a forty-third embodiment of the present invention there is provided use of bicarbonate, magnesium cations and a pH adjusting agent of the first or eighteenth embodiment or a combination of the nineteenth embodiment for the manufacture of a medicament to increase longevity in a mammal.

According to a forty-fourth embodiment of the present invention there is provided use of bicarbonate, magnesium cations and a pH adjusting agent of the first or eighteenth embodiment or a combination of the nineteenth embodiment for the manufacture of a medicament to scavenge protons in a mammal.



According to a forty-fifth embodiment of the present invention there is provided use of bicarbonate, magnesium cations and a pH adjusting agent of the first or eighteenth embodiment or a combination of the nineteenth embodiment for the manufacture of a medicament to decrease proton concentrations in a mammal.

5 According to a forty-sixth embodiment of the present invention there is provided use of bicarbonate, magnesium cations and a pH adjusting agent of the first or eighteenth embodiment or a combination of the nineteenth embodiment for the manufacture of a medicament to decrease inflammation in a mammal.

10 According to a forty-seventh embodiment of the present invention there is provided use of bicarbonate, magnesium cations and a pH adjusting agent of the first or eighteenth embodiment or a combination of the nineteenth embodiment for the manufacture of a medicament to decrease inflammatory conditions in a mammal.

15 According to a forty-eighth embodiment of the present invention there is provided use of bicarbonate, magnesium cations and a pH adjusting agent of the first or eighteenth embodiment or a combination of the nineteenth embodiment for the manufacture of a medicament to increase motor activity in a mammal.

20 According to a forty-ninth embodiment of the present invention there is provided use of bicarbonate, magnesium cations and a pH adjusting agent of the first or eighteenth embodiment or a combination of the nineteenth embodiment for the manufacture of a medicament to decrease fatigue in a mammal.

25 According to a fiftieth embodiment of the present invention there is provided an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the second embodiment or a combination of the nineteenth embodiment when used to prevent certain inflammatory diseases in a mammal in need of such prevention.

30 According to a fifty-first embodiment of the present invention there is provided an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the third embodiment or a combination of the nineteenth embodiment when used to treat certain inflammatory diseases in a mammal in need of such treatment.

35 According to a fifty-second embodiment of the present invention there is provided an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the fourth embodiment or a combination of the nineteenth embodiment when used to prevent degenerative diseases in a mammal in need of such prevention.



According to a fifty-third embodiment of the present invention there is provided an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the fifth embodiment or a combination of the nineteenth embodiment when used to treat degenerative diseases in a mammal in need of such treatment.

According to a fifty-fourth embodiment of the present invention there is provided an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the sixth embodiment or a combination of the nineteenth embodiment when used to prevent certain viral diseases in a mammal in need of such prevention.

According to a fifty-fifth embodiment of the present invention there is provided an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the seventh embodiment or a combination of the nineteenth embodiment when used to treat certain viral diseases in a mammal in need of such treatment.

According to a fifty-sixth embodiment of the present invention there is provided an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the eighth embodiment or a combination of the nineteenth embodiment when used to decrease senescence in a mammal.

According to a fifty-seventh embodiment of the present invention there is provided an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the ninth embodiment or a combination of the nineteenth embodiment when used to treat senescence in a mammal.

According to a fifty-eighth embodiment of the present invention there is provided an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the tenth embodiment or a combination of the nineteenth embodiment when used to increase longevity in a mammal.

According to a fifty-ninth embodiment of the present invention there is provided an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the eleventh embodiment or a combination of the nineteenth embodiment when used to scavenge protons in a mammal.

According to a sixtieth embodiment of the present invention there is provided an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the twelfth embodiment or a combination of the nineteenth embodiment when used to decrease proton concentrations in a mammal.



According to a sixty-first embodiment of the present invention there is provided an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the thirteenth embodiment or a combination of the nineteenth embodiment when used to decrease inflammation in a mammal.

5 According to a sixty-second embodiment of the present invention there is provided an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the fourteenth embodiment or a combination of the nineteenth embodiment when used to decrease inflammatory conditions in a mammal.

10 According to a sixty-third embodiment of the present invention there is provided an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the fifteenth embodiment or a combination of the nineteenth embodiment when used to increase motor activity in a mammal.

15 According to a sixty-fourth embodiment of the present invention there is provided an aqueous neutral to mildly alkaline metal bicarbonate solution of the first or eighteenth embodiment or a solution of the sixteenth embodiment or a combination of the nineteenth embodiment when used to decrease fatigue in a mammal.

The term mammal as used herein includes vertebrate. Examples of mammals and vertebrates to which the methods of the invention apply include a bovine, human (male or female), ovine, equine, caprine, Leporine, feline or canine mammal or vertebrate.

20 Specific examples of animals include sheep, cattle, horses, rabbits, cats, goats, alpacas, cats, dogs, pigs, rabbits, fowls, deer, buffaloes and other livestock and domestic animals.

Metal Bicarbonate Generally

Generally the pH of the aqueous metal bicarbonate solution is neutral to mildly alkaline, typically mildly alkaline and more typically in the range of 7 to 9 even more

25 typically 8 to 8.6 and the temperature of the aqueous neutral to mildly alkaline metal bicarbonate solution is maintained at such a level so as to maintain the metal bicarbonate in the aqueous diluent. The aqueous neutral to mildly alkaline metal bicarbonate solution may be kept under an atmosphere comprising carbon dioxide of from about 0.8 to 5 or 1 to 5 atmospheres, more typically 1 to 3 atmospheres and even more typically slightly

30 above atmospheric pressure such as the sorts of pressures that soft drinks are currently under in cans or bottles, for example, so as to maintain the metal bicarbonate in the aqueous diluent.



Generally the metal cation is an alkaline earth metal cation or an alkali metal cation. Generally a metal cation is chosen which is capable of acting as a bicarbonate transporter into mammalian cells. More particularly the metal cation may be cations of magnesium, sodium, potassium, calcium, lithium or any mixture thereof. Where a mixture of alkaline
5 earth metal cations or alkali metal cations are used: (1) two different alkaline earth metal



cations or alkali metal cations or mixtures thereof, the molar ratio of the first metal cation to the second may be in the range 0.5:99.5 to 99.5:0.5, typically 75:25 to 25:75, more typically 0.7:1 to 1:0.7; (2) three different alkaline earth metal cations or alkali metal cations or mixtures thereof, the molar ratio of the first metal cation to the second to the third may be in the range 99.5:0.5:0.5 to 0.5:99.5:99.5, typically 75:25:25 to 25:75:75, more typically 0.5:1:1 to 1:0.5:0.5; (3) four different alkaline earth metal cations or alkali metal cations or mixtures thereof, the molar ratio of the first metal cation to the second to the third to the fourth may be in the range 99.5:0.5:0.5:0.5 to 0.5:99.5:99.5:99.5, typically 75:25:25:25 to 25:75:75:75, more typically 0.5:1:1:1 to 0.5:1:1:1. Generally the metal cation is magnesium or a mixture of magnesium and sodium metal cations. Typically the aqueous neutral to mildly alkaline metal bicarbonate solution has a high metal cation concentration in association with bicarbonate anions.

Typically the metal bicarbonate is used at a concentration of 10-100 mole% or weight% of its saturation solubility (which will depend on the actual metal bicarbonate(s) used), more typically 10-90%, 10-80%, 10-70%, 10-60%, 10-50%, 10-40%, 10-30%, 10-20%, more typically 15-95%, 15-85%, 15-75%, 15-65%, 15-55%, 15-45%, 15-35%, 15-25%, more typically 15-90%, 15-80%, 15-70%, 15-60%, 15-50%, 15-40%, 15-30%, 15-20%, more typically 10-95%, 10-85%, 10-75%, 10-65%, 10-55%, 10-45%, 10-35%, 10-25%, more typically 20-90%, 20-80%, 20-70%, 20-60%, 20-50%, 20-40%, 20-30%, more typically 25-95%, 25-85%, 25-75%, 25-65%, 25-55%, 25-45%, 25-35%, more typically 25-90%, 25-80%, 25-70%, 25-60%, 25-50%, 25-40%, 25-30%, more typically 20-95%, 20-85%, 20-75%, 20-65%, 20-55%, 20-45%, 20-35%, more typically 30-90%, 30-80%, 30-70%, 30-60%, 30-50%, 30-40%, more typically 35-95%, 35-85%, 35-75%, 35-65%, 35-55%, 35-45%, more typically 35-90%, 35-80%, 35-70%, 35-60%, 35-50%, 35-40%, more typically 30-95%, 30-85%, 30-75%, 30-65%, 30-55%, 30-45%, more typically 40-90%, 40-80%, 40-70%, 40-60%, 40-50%, more typically 45-95%, 45-85%, 45-75%, 45-65%, 45-55%, more typically 45-90%, 45-80%, 45-70%, 45-60%, 45-50%, more typically 40-95%, 40-85%, 40-75%, 40-65%, 40-55%, more typically 50-90%, 50-80%, 50-70%, 50-60%, more typically 55-95%, 55-85%, 55-75%, 55-65%, more typically 55-90%, 55-80%, 55-70%, 55-60%, more typically 50-95%, 50-85%, 50-75%, 50-65%, more typically 60-90%, 60-80%, 60-70%, more typically 65-95%, 65-85%, 65-75%, more typically 65-90%, 65-80%, 65-70%, more typically 60-95%, 60-85%, 60-75%, more typically 70-90%, 70-80%, more typically 75-95%, 75-85%, more typically 75-90%, 75-

80%, more typically 70-95%, 70-85%, more typically 80-90%, more typically 85-95%, more typically 85-90%, more typically 80-95%, more typically 20-100%, 30-100%, 40-100%, 50-100%, 60-100%, 70-100%, 80-100% or 90-100%. Depending on the solubility of the metal bicarbonate, the amount of metal cation may range from 20mg to 1250mg or 25mg to 1250mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, typically 20mg to 1000mg or 50mg to 1000mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, more typically 20mg to 750mg or 50mg to 750mg or 20mg to 600mg or 50mg to 600mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, even more typically 20mg to 500mg or 30mg to 500mg or 50mg to 500mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, even more typically 20mg to 250mg or 50mg to 250mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, most typically 100mg to 500mg or 100mg to 400mg or 100mg to 300mg or 100mg to 250mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, even most typically 20mg to 200mg or 20 to 150mg or 20mg to 120mg or 120mg to 300mg or 120mg to 200mg. Typically when the metal cation is magnesium, the amount of magnesium may range from 30mg to 140mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, typically 30mg to 130mg, 30mg to 120mg, 30mg to 110mg, 30mg to 100mg, 30mg to 90mg, 30mg to 80mg, 30mg to 70mg, 30mg to 60mg, 30mg to 50mg, 30mg to 40mg, 50mg to 120mg, 60mg to 120mg, 70mg to 120mg, 80mg to 120mg, 90mg to 120mg or 75mg to 120mg or 100mg to 120mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution. Typically when the metal cation is sodium and/or potassium, the amount of sodium and/or potassium may range from greater than 30mg to 1250mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, typically 50mg to 1000mg or 50mg to 750mg or 50mg to 500mg or 75mg to 1250mg or 75mg to 1000mg or 75mg to 500mg or 100mg to 1000mg or 100mg to 500mg or 250mg to 1000mg or 250mg to 500mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution. Typically when the metal cation is calcium, the amount of calcium may range from greater than 20mg to 1250mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, typically 20mg to 1000mg or 20mg to 750mg or 20mg to 500mg or 20mg to 250mg or 20mg to 200mg or 20mg to 150mg or 20mg to 100mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution. Typically the amount of bicarbonate anion present will be stoichiometric with the amount of metal cation in solution so as to form the metal

bicarbonate. Alternatively, the concentration of the metal bicarbonate can be based on the bicarbonate anion concentrations in which case the amount of bicarbonate anion (which will depend on the saturation solubility of the actual metal bicarbonate anion(s) used).

The concentration of bicarbonate typically ranges from 120mg or 150mg to 3500mg per 5 litre of aqueous neutral to mildly alkaline metal bicarbonate solution, typically 120mg or 150mg to 3000mg or 200mg to 3000mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, more typically 250mg to 2100mg or 300mg to 2000mg or 200mg to 1500mg or 300mg to 1500mg or 400mg to 1500mg or 500mg to 1500mg or 600mg to 1500mg or 700mg to 1500mg or 800mg to 1500mg or 900mg to 1500mg or 10 1000mg to 1500mg or 200mg to 1000mg or 300mg to 1000mg or 400mg to 1000mg or 500mg to 1000mg or 600mg to 1000mg or 700mg to 1000mg or 800mg to 1000mg or 900mg to 1000mg or 1000mg to 1500mg or 1200mg to 1500mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, even more typically 600mg to 1000mg or 500mg to 1500mg per litre of aqueous neutral to mildly alkaline metal
15 bicarbonate solution, most typically 950mg or 200mg to 2000mg or 200mg to 1750mg or 200mg to 1250mg or 200mg to 100mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution. Typically at least 600mg of bicarbonate anions per litre of solution is present, more typically 600-1800mg/l, 600-1500mg/l, 600-1350mg/l, 600-1200mg/l, 600-1100mg/l, 600-1000mg/l, 600-950mg/l, 600-900mg/l, 600-850mg/l, 600-800mg/l,
20 600-750mg/l, 600-700mg/l or 600-650mg/l. Typically a mildly alkaline saturated magnesium bicarbonate solution is used or a mildly alkaline solution comprising a mixture of sodium and/or potassium and magnesium bicarbonate, more typically sodium and magnesium bicarbonate. Typically the range for a mixture of sodium and/or potassium and magnesium bicarbonate, more typically sodium and magnesium bicarbonate varies
25 from 20mg to 1250mg or 25mg to 1250mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, typically 20mg to 1000mg or 50mg to 1000mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, more typically 20mg to 750mg or 50mg to 750mg or 20mg to 600mg or 50mg to 600mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, even more typically 20mg to 500mg
30 or 30mg to 500mg or 50mg to 500mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, even more typically 20mg to 250mg or 50mg to 250mg or even more typically 20mg to 300mg or 50mg to 300mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, most typically 75mg to 1000mg or 75mg to 500mg or

100mg to 1000mg or 100mg to 500mg or 100mg to 400mg or 100mg to 300mg or 100mg to 250mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, even most typically 20mg to 200mg or 20 to 150mg or 20mg to 120mg or 120mg to 300mg or 120mg to 200mg. Usually the ratio (weight to weight) of magnesium to sodium is in the range 25:1 to 1:4, typically 1:1.125.

Generally the aqueous diluent is water or comprises water. Generally the carbon dioxide-containing-aqueous diluent may be carbonic acid in water, hydrated carbon dioxide in water, carbon dioxide gas dissolved in water, carbonated soft drinks, carbonated mineral water, soda water or other carbon dioxide-containing-aqueous diluents. If carbon dioxide gas is used, the carbon dioxide may be either bubbled into aqueous solutions containing metal carbonate or metal carbonate hydroxide or metal oxide or mixture thereof or the carbon dioxide may be introduced in the form of a blanket over aqueous solutions containing metal carbonate or metal carbonate hydroxide or metal oxide or mixture thereof. Typically the carbon dioxide-containing-aqueous diluent is pharmaceutically acceptable. Typically carbonated mineral water, carbonic acid, hydrated carbon dioxide in water or carbonated water is used. The amounts of carbon dioxide-containing-aqueous diluent and metal carbonate or metal carbonate hydroxide or metal oxide or mixture thereof used are sufficient to obtain a clear solution at a neutral to mildly alkaline pH, typically pH 7 to 9 or pH 7 to 8.6, more typically pH 7.5 to 8.8 or pH 7.5 to 8.5 or pH 7.8 to 8.6, pH 7.8 to 8.5, pH 7.8 to 8.4, pH 7.8 to 8.3, pH 7.8 to 8.2, pH 7.8 to 8.1, pH 7.8 to 8.0, pH 7.8 to 7.9, pH 7.9 to 8.6, pH 7.9 to 8.5, pH 7.9 to 8.4, pH 7.9 to 8.3, pH 7.9 to 8.2, pH 7.9 to 8.1, pH 7.9 to 8.0, pH 8.0 to 8.6, pH 8.0 to 8.5, pH 8.0 to 8.4, pH 8.0 to 8.3, pH 8.0 to 8.2, pH 8.0 to 8.1, pH 8.1 to 8.6, pH 8.1 to 8.5, pH 8.1 to 8.4, pH 8.1 to 8.3, pH 8.1 to 8.2, pH 8.2 to 8.6, pH 8.2 to 8.5, pH 8.2 to 8.4, pH 8.2 to 8.3, pH 8.3 to 8.6, pH 8.3 to 8.5, pH 8.3 to 8.4, pH 8.4 to 8.6, pH 8.4 to 8.5, pH 8.5 to 8.6, even more typically pH 8 to 8.5 or pH 8.2 to 8.6, most typically pH 8.3. Usually 10 to 60mL, typically 25 to 55mL, more typically 40 to 50mL, most typically approximately 45mL of chilled carbonated mineral water per litre of water is used. Usually the chilled carbonated mineral water is at a temperature of 0 to 25°C, 0 to 20°C, 0.5 to 25°C, 0.5 to 20°C, 0.5 to 15°C, 0.5 to 10°C, 0.5 to 9°C, 0.5 to 8°C, 0.5 to 7°C, 1 to 20°C, 1 to 15°C, 1 to 10°C, 1.5 to 20°C, 1.5 to 15°C, 1.5 to 10°C, 2 to 20°C, 2 to 15°C, 2 to 10°C, 3 to 20°C, 3 to 15°C, 4 to 20°C, 4 to 15°C, 4 to 10°C, 5 to 20°C, 5 to 15°C, 6 to 20°C, 6 to 15°C, 6 to 10°C, 7 to 20°C, 7 to 15°C, 7 to 10°C, 8 to 20°C, 8 to 15°C, 8 to 10°C, 9 to

even more typically 3°C to 10°C, most typically 5°C to 10°C and even most typically 5°C.

Generally the pH adjusting agent is carbon dioxide gas, carbonic acid in water, hydrated carbon dioxide in water, carbon dioxide gas in water, carbonated soft drinks, carbonated mineral water, soda water or other carbon dioxide-containing-aqueous diluents or an alkali or any mixture thereof. Examples of alkalis are water soluble drinkable alkalis such as sodium hydroxide, sodium carbonate, potassium carbonate or potassium hydroxide or any mixture thereof.

Typically additives may be added during the process of the invention or to the aqueous neutral to mildly alkaline metal bicarbonate solution. The additives may be 0mg or 0.5mg to 1000mg sodium bicarbonate per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, typically 25mg to 900mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, typically 50mg to 800mg or 50mg to 500mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, more typically 100mg to 700mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, even more typically 200mg to 600mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, most typically 300mg to 500mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, even most typically 500mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution. The additives may also be chlorides and other appropriate salts of magnesium, sodium, potassium, calcium and lithium, such as carbonates or hydroxides or sulfates, with or without the addition of sodium bicarbonate. For example, magnesium sulfate, magnesium chloride or other soluble salts of magnesium. Further additives may include potassium bicarbonate, calcium bicarbonate or lithium bicarbonate. Generally calcium bicarbonate is prepared by adding carbonic acid or carbonated water or hydrated carbon dioxide or carbon dioxide gas to a mixture of calcium carbonate in water. Generally lithium bicarbonate is prepared by adding carbonic acid and/or carbonated water and/or hydrated carbon dioxide and/or carbon dioxide gas and/or solid carbon dioxide to a mixture of lithium carbonate in water.

The aqueous neutral to mildly alkaline metal bicarbonate solution may further include a stabilising agent. The stabilising agent may also be a pH adjusting agent. Typically the stabilising agent is a gaseous phase, for example carbon dioxide gas, which maintains

20°C, 9 to 15°C, 9 to 10°C, 10 to 15°C, typically 0 to 15°C, more typically 0 to 10°C, even more typically 3°C to 10°C, most typically 5°C to 10°C and even most typically 5°C. Alternatively the metal carbonate or metal carbonate hydroxide or metal oxide or mixture thereof can be added after the carbon dioxide has been added.

- 5 Generally the metal bicarbonate in aqueous solution may be derived from a metal carbonate or metal carbonate hydroxide or metal oxide or metal bicarbonate or metal hydroxide or other appropriate metal compound or any mixture thereof. Examples include magnesium, sodium, potassium, calcium, lithium carbonate or carbonate hydroxide or oxide or bicarbonate or a mixture of any two or more thereof. For example
- 10 magnesium carbonate hydroxide pentahydrate, the calcite series or dolomite series of minerals (Mg, Ca)CO₃ or limestone or dolomite rocks is used. Generally magnesium carbonate hydroxide pentahydrate or a mixture of magnesium carbonate hydroxide pentahydrate and sodium bicarbonate is used.

Generally the pH of the aqueous metal bicarbonate solution for oral administration is

15 neutral to mildly alkaline, typically in the range pH 7 to 9 or pH 7 to 8.6, more typically pH 7.5 to 8.8 or pH 7.5 to 8.5 or pH 7.8 to 8.6, pH 7.8 to 8.5, pH 7.8 to 8.4, pH 7.8 to 8.3, pH 7.8 to 8.2, pH 7.8 to 8.1, pH 7.8 to 8.0, pH 7.8 to 7.9, pH 7.9 to 8.6, pH 7.9 to 8.5, pH 7.9 to 8.4, pH 7.9 to 8.3, pH 7.9 to 8.2, pH 7.9 to 8.1, pH 7.9 to 8.0, pH 8.0 to 8.6, pH 8.0 to 8.5, pH 8.0 to 8.4, pH 8.0 to 8.3, pH 8.0 to 8.2, pH 8.0 to 8.1, pH 8.1 to

20 8.6, pH 8.1 to 8.5, pH 8.1 to 8.4, pH 8.1 to 8.3, pH 8.1 to 8.2, pH 8.2 to 8.6, pH 8.2 to 8.5, pH 8.2 to 8.4, pH 8.2 to 8.3, pH 8.3 to 8.6, pH 8.3 to 8.5, pH 8.3 to 8.4, pH 8.4 to 8.6, pH 8.4 to 8.5, pH 8.5 to 8.6, even more typically pH 8 to 8.5 or pH 8.2 to 8.6, most typically pH 8.3. Generally the pH of the aqueous metal bicarbonate solution for parenteral administration is neutral to very mildly alkaline, typically in the range pH 7 to

25 7.6, or pH 7.0 to 7.5, or pH 7.1 to 7.5, more typically pH 7.2 to 7.5 or pH 7.3 to 7.5 or pH 7.4 to 7.5. Generally the aqueous neutral to mildly alkaline metal bicarbonate solution is prepared and stored at a temperature ranging from 0 to 25°C, 0 to 20°C, 0.5 to 25°C, 0.5 to 20°C, 0.5 to 15°C, 0.5 to 10°C, 0.5 to 9°C, 0.5 to 8°C, 0.5 to 7°C, 1 to 20°C, 1 to 15°C, 1 to 10°C, 1.5 to 20°C, 1.5 to 15°C, 1.5 to 10°C, 2 to 20°C, 2 to 15°C, 2 to 10°C,

30 3 to 20°C, 3 to 15°C, 4 to 20°C, 4 to 15°C, 4 to 10°C, 5 to 20°C, 5 to 15°C, 6 to 20°C, 6 to 15°C, 6 to 10°C, 7 to 20°C, 7 to 15°C, 7 to 10°C, 8 to 20°C, 8 to 15°C, 8 to 10°C, 9 to 20°C, 9 to 15°C, 9 to 10°C, 10 to 15°C, typically 0 to 15°C, more typically 0 to 10°C,

and/or stabilises the solution at a pH of 7 to 9 and at a temperature of 0 to 55°C more typically 0 to 25°C.

Generally once the solution is prepared, the solution may be stored under a blanket of carbon dioxide gas or a mixture of carbon dioxide gas and a nondeleterious inert gas, for example, argon, helium, air, oxygen and/or nitrogen wherein the amount of carbon dioxide present in the inert gas is sufficient to maintain the solution at a pH of 7 to 9 and at a temperature of 0 to 25°C and to prevent the metal bicarbonate from forming insoluble compounds which can precipitate out of solution. Typically the carbon dioxide gas above the solution prevents loss of carbon dioxide from the solution. The amount of carbon dioxide in the gaseous mixture provides partial pressure on the liquid which is substantially equal to the partial pressure which is produced from equilibrium of bicarbonate in the solution at the mixing temperature.

Magnesium Bicarbonate Particularly

Typically the production of magnesium bicarbonate utilises the dissolution of magnesium carbonate by carbonic acid or hydrated carbon dioxide solutions. Ideally, the dissolution is produced within a defined range of conditions - a defined range of pH values, a defined range of temperature values and a defined minimum time. For optimal biological and medical activities, and for therapeutic safety, the concentrations of the component ions are defined also.

Typically to prepare the aqueous neutral to mildly alkaline metal bicarbonate solution, crushed or powdered metal carbonate, or metal carbonate hydroxide or metal oxide, such as magnesium carbonate MgCO_3 , or commercial magnesium carbonate hydroxide pentahydrate $(\text{MgCO}_3)_4 \cdot \text{Mg}(\text{OH})_2 \cdot 5\text{H}_2\text{O}$, or other commercial magnesium carbonate hydroxides, or hydrated magnesium oxides, or magnesium oxides heated with carbon dioxide, or the calcite series or dolomite series of minerals $(\text{Mg, Ca})\text{CO}_3$, or limestone or dolomite rocks is mixed with water. A cloudy suspension is obtained. Sufficient carbonic acid and/or hydrated carbon dioxide and/or carbon dioxide gas and/or solid carbon dioxide is added to obtain a solution having a pH 7 to 9 or pH 7 to 8.6, more typically pH 7.5 to 8.8 or pH 7.5 to 8.5 or pH 7.8 to 8.6, pH 7.8 to 8.5, pH 7.8 to 8.4, pH 7.8 to 8.3, pH 7.8 to 8.2, pH 7.8 to 8.1, pH 7.8 to 8.0, pH 7.8 to 7.9, pH 7.9 to 8.6, pH 7.9 to 8.5, pH 7.9 to 8.4, pH 7.9 to 8.3, pH 7.9 to 8.2, pH 7.9 to 8.1, pH 7.9 to 8.0, pH 8.0 to 8.6, pH 8.0 to 8.5, pH 8.0 to 8.4, pH 8.0 to 8.3, pH 8.0 to 8.2, pH 8.0 to 8.1, pH 8.1 to 8.6,

pH 8.1 to 8.5, pH 8.1 to 8.4, pH 8.1 to 8.3, pH 8.1 to 8.2, pH 8.2 to 8.6, pH 8.2 to 8.5, pH 8.2 to 8.4, pH 8.2 to 8.3, pH 8.3 to 8.6, pH 8.3 to 8.5, pH 8.3 to 8.4, pH 8.4 to 8.6, pH 8.4 to 8.5, pH 8.5 to 8.6, even more typically pH 8 to 8.6 or pH 8.2 to 8.6, most typically pH 8.3. The solution is then typically placed in a closed or sealed container at 0 to 20°C or 0 to 15°C with occasional mixing until a clear solution develops. The amount of carbonic acid and/or hydrated carbon dioxide and/or carbon dioxide gas bubbled through the solution and dissolved therein and/or solid carbon dioxide is sufficient to prevent precipitation of water insoluble metal compounds (such as magnesium or calcium carbonate). A clear solution is generally obtained in about 6 hours to 7 days, typically 12 to 24 hours to 5 days, more typically 24 hours to 5 days, most typically 24 hours to 3 days. Generally the aqueous neutral to mildly alkaline metal bicarbonate solution is prepared and stored at a temperature ranging from 0 to 55°C, 0 to 25°C, 0 to 20°C, 0.5 to 25°C, 0.5 to 20°C, 0.5 to 15°C, 0.5 to 10°C, 0.5 to 9°C, 0.5 to 8°C, 0.5 to 7°C, 1 to 20°C, 1 to 15°C, 1 to 10°C, 1.5 to 20°C, 1.5 to 15°C, 1.5 to 10°C, 2 to 20°C, 2 to 15°C, 2 to 10°C, 3 to 20°C, 3 to 15°C, 4 to 20°C, 4 to 15°C, 4 to 10°C, 5 to 20°C, 5 to 15°C, 6 to 20°C, 6 to 15°C, 6 to 10°C, 7 to 20°C, 7 to 15°C, 7 to 10°C, 8 to 20°C, 8 to 15°C, 8 to 10°C, 9 to 20°C, 9 to 15°C, 9 to 10°C, 10 to 15°C, typically 0 to 15°C, more typically 0 to 10°C, even more typically 3°C to 10°C, most typically 5°C to 10°C and even most typically 5°C. Alternatively the crushed or powdered metal carbonate, or metal carbonate hydroxide or metal oxide or mixture thereof is added to an aqueous solution of the carbonic acid and/or hydrated carbon dioxide and/or to an aqueous solution through which carbon dioxide gas is bubbled and/or solid carbon dioxide has been added. The amount of carbonic acid and/or hydrated carbon dioxide and/or carbon dioxide gas bubbled through the solution and dissolved therein and/or solid carbon dioxide is sufficient to prevent precipitation of water insoluble metal compounds (such as magnesium or calcium carbonate).

Typically one litre of water is placed in a container and sufficient carbonic acid and/or carbonated water and/or hydrated carbon dioxide and/or carbon dioxide gas and/or solid carbon dioxide is added to produce a pH value of approximately pH 5.2. (In practice, approximately 40 to 45mL of chilled (5°C) carbonated mineral water is used depending on the initial pH of the water). The container is sealed and the contents are mixed. 485mg magnesium carbonate hydroxide pentahydrate powder ($\text{MgCO}_3 \cdot 4\text{Mg}(\text{OH})_2 \cdot 5\text{H}_2\text{O}$, molecular weight 485) is added. The container is again sealed and the contents are mixed.

The container is stored at a temperature of 0 to 10°C and the contents mixed regularly. Sufficient time is allowed for a clear solution of magnesium bicarbonate to develop at a range of pH 8.0 to pH 8.6, preferably pH 8.3. This takes approximately 24 to 72 hours. Alternatively the carbonic acid and/or carbonated water and/or hydrated carbon dioxide and/or carbon dioxide gas and/or solid carbon dioxide is added to the magnesium carbonate hydroxide pentahydrate powder in water. Alternatively one litre of water is placed in a container and sufficient carbonic acid and/or carbonated water and/or hydrated carbon dioxide and/or solid carbon dioxide is added to produce a pH value less than pH 5.2. (In practice, approximately 30mL to 40mL of chilled water is used depending on the initial pH of the water). The container is sealed and the contents are mixed. 485mg magnesium carbonate hydroxide pentahydrate powder $(\text{MgCO}_3)_4 \cdot \text{Mg}(\text{OH})_2 \cdot 5\text{H}_2\text{O}$, molecular weight 485 is added. The container is again sealed and the contents are mixed. The container is stored at a temperature of 0 to 10°C and the contents mixed regularly. The pH of the water is then adjusted with an alkali such as sodium hydroxide or potassium hydroxide to a pH of 8 to 8.6, typically pH 8.3. Alternatively the carbonic acid or carbonated water and/or hydrated carbon dioxide and/or carbon dioxide gas and/or solid carbon dioxide is added to the magnesium carbonate hydroxide pentahydrate powder in water.

The above processes may optionally be conducted under an atmosphere of carbon dioxide or a gas comprising carbon dioxide.

Generally once the solution is prepared, it may be stored under a blanket of carbon dioxide gas to maintain the solution at a pH of 7 to 9 and at a temperature of 0 to 25°C.

Usually one litre of the magnesium bicarbonate solution prepared above contains approximately 120mg of magnesium per litre of aqueous neutral to mildly alkaline metal bicarbonate solution and approximately 600mg of bicarbonate. 500mg sodium bicarbonate (or potassium bicarbonate) is added to the magnesium bicarbonate solution and mixed. The mixture is stored in a sealed container in a refrigerator. The mixture contains approximately 120mg magnesium per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, 135mg sodium per litre of aqueous neutral to mildly alkaline metal bicarbonate solution and 950mg bicarbonate per litre of aqueous neutral to mildly alkaline metal bicarbonate solution.

Generally the aqueous neutral to mildly alkaline metal bicarbonate solution of the invention is administered or consumed orally. Typically the solution is an orally drinkable solution. Typically the solution is a therapeutic orally drinkable solution. Alternatively a gelling agent may added to the solution and the solution subjected to gelling conditions to gel the solution and the resultant gel may be consumed orally. For example, the aqueous neutral to mildly alkaline metal bicarbonate solution may be prepared as a solution or an iced confectionary, such as an ice block or iced dessert, which is ingested orally. Alternatively the aqueous neutral to mildly alkaline metal bicarbonate solution may be prepared in the form of a tablet, lozenge or lolly which is ingested orally. For example, the aqueous neutral to mildly alkaline metal bicarbonate solution may be administered for metabolic acidosis or renal failure. Optionally the solution may be sterilised. Typically the aqueous neutral to mildly alkaline metal bicarbonate solution is prepared as a solution which is ingested on a regular basis hourly, daily, monthly or yearly. The amount and frequency of aqueous neutral to mildly alkaline metal bicarbonate solution administered/consumed in a day is generally sufficient so as to maintain a steady bicarbonate level in the bicarbonate concentration of a taker's body fluids. It is preferable to avoid a rapid increase in the bicarbonate level in the bicarbonate concentration of a taker's body fluids. The amount of aqueous neutral to mildly alkaline metal bicarbonate solution administered in a day ranges from 250mL to 6 litres, typically 250mL to 5.5 litres, 250mL to 5 litres, 250mL to 4.5 litres, 250mL to 4 litres, 250mL to 3.5 litres, 250mL to 3 litres, 500mL to 6 litres, 500mL to 5.5 litres, 500mL to 5 litres, 500mL to 4.5 litres, 500mL to 4 litres, 500mL to 3.5 litres, 500mL to 3 litres, more typically 1 litre to 6 litres, 1 litre to 5.5 litres, 1 litre to 5 litres, 1 litre to 4.5 litres, 1 litre to 4 litres, 1 litre to 3.5 litres, even more typically 1 litre to 3 litres, 1.5 litres to 6 litres, 1.5 litres to 5.5 litres, 1.5 litres to 5 litres, 1.5 litres to 4.5 litres, 1.5 litres to 4 litres, 1.5 litres to 3.5 litres, 1.8 litres to 3.3 litres, 1.8 to 2.8 litres, 1.8 to 2.5 litres, 1.8 to 2.3 litres, 1.8 to 2.0 litres, most typically 2 to 3 litres, typically 2.3 to 2.8 litres, more typically 2.3 to 2.6 litres, usually 2.1 to 3 litres. The aqueous neutral to mildly alkaline metal bicarbonate solution may be administered on a full or empty stomach, typically the aqueous neutral to mildly alkaline metal bicarbonate solution is administered on an empty stomach. Usually 1.5 to 3.5 litres, typically 1.8 to 3 litres, more typically 1.5 to 2.4 litres, even more typically 1.8 to 2.1 litres and usually between 1.8 and 2.7 litres of aqueous neutral to mildly alkaline metal bicarbonate solution is ingested, administered or

consumed on an empty stomach by a mammal (typically a human) in equal or non equal volume amounts (100mL - 1000mL, 200-800mL, 250-750mL, 275-700mL, 300-650mL, 350-600mL, 400-550mL, 450-500mL, typically about 300-400mL, more typically about 375mL volume amounts a number of times (typically at set times) each day for the required number of times per day to drink the desired daily amount of the solution). For example if 1800mL per day is to be consumed then a user may drink six 300mL amounts of the solution every 2 to 2.5 hours throughout the day. The oral consumption of the solution three or more times at roughly equally spaced apart intervals throughout the day is more desirable than consuming the solution in one or two lots throughout the day. The idea of taking the solution is to take it regularly throughout the day so that a simulated continuous oral intake or a close to continuous regular oral intake of the solution occurs. Thus depending on the condition and the subject one suitable administration/consumption regime could be nine by 200mL amounts of the solution, each 200mL amount being orally administered/consumed about every 1.5-1.75 hours to provide a total daily intake of 1800mL. Alternatively, once again depending on the condition and the subject one suitable administration/consumption regime could be nine by 300mL amounts of the solution, each 300mL amount being orally administered/consumed about every 1.5-1.75 hours to provide a total daily intake of 2700mL. Alternatively, once again depending on the condition and the subject one suitable administration/consumption regime could be nine by 350mL amounts of the solution, each 350mL amount being orally administered/consumed about every 1.5-1.75 hours to provide a total daily intake of 3150mL. Typically the solution is administered/consumed 3 to 30, 3-25, 3-20, 3-15, 3-12, 3-10, 3-9, 3-8, 3-7, 3-6, 3-5, 3-4, 4-30, 4-25, 4-20, 4-15, 4-12, 4-10, 4-9, 4-8, 4-7, 4-6, 4-5, 5-30, 5-25, 5-20, 5-15, 5-12, 5-11, 5-10, 5-9, 5-8, 5-7, 5-6, 6-30, 6-25, 6-20, 6-15, 6-12, 6-11, 6-10, 6-9, 6-8, 6-7, 7-30, 7-25, 7-20, 7-15, 7-12, 7-11, 7-10, 7-9, 7-8, 8-30, 8-25, 8-20, 8-15, 8-12, 8-11, 8-10, 8-9 times per day at regular or irregular intervals or a mixture of both regular and irregular intervals, throughout each day. Typically the solution is administered/consumed every .3-10, .3-8, .3-7, .3-6, .3-5, .3-4.5, .3-4, .3-3.5, .3-3, .3-2.5, .3-2, .3-1.5, .3-1, .3-.75, .3-.5 hours/day when the subject is awake. More typically the solution is administered/consumed every 0.5-8, 0.5-7, 0.5-6, .5-5, .5-4.5, .5-4, .5-3.5, .5-3, .5-2.5, .5-2, .5-1.5, .5-1, .5-.8, .5-.75 hours/day when the subject is awake. Where possible the solution is consumed/administered on an empty (e.g. before eating). The solution may be administered according to these latter dosages over short (for

example 1 to 60 days, 10 to 40 days, 3 months to 6 months, 1 day to 6 months) or long (for example 6 months to 10 years or more, 9 months to 18 months, 1 year to 3 years, 1 year to 5 years, 2 to 6 years) periods as required. Usually the amount of aqueous neutral to mildly alkaline metal bicarbonate solution administered to a mammal is 5 to 100mL per Kg, more 5 usual 10 to 50mL per Kg, most usual 14 to 29mL per Kg or 25 to 43mL per Kg.

The solution may include other additives such as sweeteners, preservatives, flavourings and other suitable additives. Examples of suitable sweeteners include sucrose, lactose, glucose, aspartame or saccharine. Examples of suitable flavouring agents include peppermint oil, oil of wintergreen, cherry, orange or raspberry flavouring. Examples of suitable preservatives 10 include sodium benzoate, vitamin E, alpha-tocopherol, ascorbic acid, methyl paraben, propyl paraben or sodium bisulphite.

Typically the aqueous neutral to mildly alkaline metal bicarbonate solution is orally administered/consumed on an empty stomach. Usually consumption in this manner avoids the mixing of bicarbonate anions with stomach acid which may result in the loss of 15 bicarbonate. Usually the aqueous neutral to mildly alkaline metal bicarbonate solution is consumed in small amounts a number of times through a day typically at set times each day to avoid a rapid increase in the bicarbonate concentration of body fluids. Usually the amount of aqueous neutral to mildly alkaline metal bicarbonate solution consumed at commencement is 500mL per day and is increased by increments over a period of one 20 month to the maximum consumption. This start-up schedule generally avoids any gastrointestinal side effects due to the smooth muscle relaxation properties of magnesium.

The aqueous neutral to very mildly alkaline metal bicarbonate solution of the invention may be administered intravenously (e.g. by discrete injection, semi continuous injection or drip feed or continuous injection or drip feed) or by other parenteral routes. Another 25 embodiment of the invention is directed to a pharmaceutical composition comprising the solution of the first or eleventh embodiments together with one or more pharmaceutically acceptable carriers, diluents, adjuvants and/or excipients. Typically the pharmaceutical composition is suitable for oral or parenteral administration. Another embodiment of the invention is directed to a veterinary composition comprising the solution of the first or 30 eleventh embodiments together with one or more veterinarily acceptable carriers, diluents, adjuvants and/or excipients. Typically the veterinary composition is suitable for oral or parenteral administration. The amount and frequency of aqueous neutral to mildly alkaline metal bicarbonate solution administered/consumed in a day is generally sufficient so as to

maintain a steady bicarbonate level in the bicarbonate concentration of a taker's body fluids. It is preferable to avoid a rapid increase in the bicarbonate level in the bicarbonate concentration of a taker's body fluids. For parenteral administration, the solution is generally sterile. Suitable mono-toxic parenterally acceptable diluents or solvents include 5 water, Ringer's solution, isotonic salt solution, 1,3-butanediol, ethanol, propylene glycol or polyethylene glycols in mixtures with water. Aqueous solutions or suspensions may further comprise one or more buffering agents. Suitable buffering agents include sodium borate, sodium acetate, sodium citrate, or sodium tartrate, for example. Typically the solution is administered on a regular basis throughout a day to a patient requiring treatment. For 10 example a patient may be parenterally administered the solution by way of a continuous drip feed or alternatively by way of a number of injections of the solution throughout a day (e.g. every .5-8 hours, more typically every 1-4 hours). The treatment is generally continued as long as required to alleviate the patient's symptoms to a satisfactory level. For concentration of metal bicarbonate in the compositions, frequency of administration and amount 15 administered see discussion under oral administration.

Brief Description of Drawings

Fig. 1 are plots of the survival curves for a control group of sheep and a treatment group of sheep;

Fig. 2 is a photograph showing osteoarthritis in the joints of the fingers and thumb. 20 Osteoarthritis before the consumption of aqueous metal bicarbonate solution. Note the swelling and 'claw-like' hand resulting from joint flexion and joint displacement. (The patient was pushing down with her hand '*as hard as possible*' in an attempt to place her hand flat on the underlying surface.); and


Fig. 3 is a photograph showing osteoarthritis in the joints of the fingers and thumb. 25 Osteoarthritis twelve months after commencement of the consumption of aqueous metal bicarbonate solution. Note that the fingers can be extended and the joints are '*straighter*' than twelve months previously. (The patient had placed her hand flat on the underlying surface without exerting any force.)

Best Mode And Other Mode(s) For Carrying Out The Invention

30 Magnesium bicarbonate is a natural hydrated salt which exists only in an aqueous solution. It may be formed in spring water by an ion exchange process between the protons in carbonic acid (formed from the hydration of carbon dioxide located in the atmosphere,

organic material, soils and rocks) and the magnesium in the constituent minerals of rocks (particularly the ferromagnesian minerals known as pyroxene and olivine that constitute basalt rocks).

The ion exchange process can be represented by the following equations:





pH dependent

- 10 The term magnesium bicarbonate is used universally to describe the mixture of magnesium cations and bicarbonate anions found in spring waters and mineral waters. Most spring waters and mineral waters have acidic pH values (about pH 6.0). If the pH value of the water rises (due to contact with hydroxides), the magnesium cations and bicarbonate anions "attract" each other (reversibly) to form the true salt. The chemical formula of magnesium bicarbonate may be written as $\text{Mg}(\text{HCO}_3)_2$, or (more accurately) $\text{Mg}(\text{H}_2\text{O})_4(\text{HCO}_3)_2$. This latter formula takes into account the hexahydrated magnesium cation $\text{Mg}(\text{H}_2\text{O})_6^{2+}$.

In essence, magnesium bicarbonate exists in aqueous solution probably as an hydrated salt of indeterminate hydration size due to the hydrogen bonds between linked water dipoles centred around the hydrated magnesium cation.

The chemical processes occurring in magnesium bicarbonate solutions are complex and depend on the concentrations of magnesium cations and other ions. The following reactions are considered to occur:





5 There exists also a range of possible acid-base equilibria involving HCO_3^- , H_3O^+ , CO_3^{2-} and OH^- ions and CO_2 and H_2CO_3 . Usually the pH adjusting agent (and/or stabilising agent) maintains the acid-base equilibria.

Typically to prepare the aqueous neutral to mildly alkaline metal bicarbonate solution, crushed or powdered metal carbonate, or metal carbonate hydroxide or metal oxide, such
 10 as magnesium carbonate MgCO_3 , or commercial magnesium carbonate hydroxide pentahydrate $(\text{MgCO}_3)_4 \cdot \text{Mg(OH)}_2 \cdot 5\text{H}_2\text{O}$, or other commercial magnesium carbonate hydroxides, or hydrated magnesium oxides, or magnesium oxides heated with carbon dioxide, or the calcite series or dolomite series of minerals $(\text{Mg, Ca})\text{CO}_3$, or limestone or dolomite rocks is mixed with water. A cloudy suspension is obtained. Sufficient carbonic
 15 acid or hydrated carbon dioxide or carbon dioxide gas is added to obtain a solution having a pH 7 to 9 or pH 7 to 8.6, more typically pH 7.5 to 8.8 or pH 7.5 to 8.5 or pH 7.8 to 8.6, pH 7.8 to 8.5, pH 7.8 to 8.4, pH 7.8 to 8.3, pH 7.8 to 8.2, pH 7.8 to 8.1, pH 7.8 to 8.0, pH 7.8 to 7.9, pH 7.9 to 8.6, pH 7.9 to 8.5, pH 7.9 to 8.4, pH 7.9 to 8.3, pH 7.9 to 8.2, pH 7.9 to 8.1, pH 7.9 to 8.0, pH 8.0 to 8.6, pH 8.0 to 8.5, pH 8.0 to 8.4, pH 8.0 to
 20 8.3, pH 8.0 to 8.2, pH 8.0 to 8.1, pH 8.1 to 8.6, pH 8.1 to 8.5, pH 8.1 to 8.4, pH 8.1 to 8.3, pH 8.1 to 8.2, pH 8.2 to 8.6, pH 8.2 to 8.5, pH 8.2 to 8.4, pH 8.2 to 8.3, pH 8.3 to 8.6, pH 8.3 to 8.5, pH 8.3 to 8.4, pH 8.4 to 8.6, pH 8.4 to 8.5, pH 8.5 to 8.6, even more typically pH 8 to 8.5 or pH 8.2 to 8.6, most typically pH 8.3. The solution is then typically placed in a sealed container at 0 to 20°C with occasional mixing until a clear
 25 solution develops. A clear solution is generally obtained in about 6 hours to 7 days, typically 12 hours to 5 days, more typically 24 hours to 5 days, most typically 24 hours to 3 days. Generally the aqueous neutral to mildly alkaline metal bicarbonate solution is prepared and stored at a temperature ranging from 0 to 25°C, 0 to 20°C, 0.5 to 25°C, 0.5 to 20°C, 0.5 to 15°C, 0.5 to 10°C, 0.5 to 9°C, 0.5 to 8°C, 0.5 to 7°C, 1 to 20°C, 1 to
 30 15°C, 1 to 10°C, 1.5 to 20°C, 1.5 to 15°C, 1.5 to 10°C, 2 to 20°C, 2 to 15°C, 2 to 10°C, 3 to 20°C, 3 to 15°C, 4 to 20°C, 4 to 15°C, 4 to 10°C, 5 to 20°C, 5 to 15°C, 6 to 20°C, 6 to 15°C, 6 to 10°C, 7 to 20°C, 7 to 15°C, 7 to 10°C, 8 to 20°C, 8 to 15°C, 8 to 10°C, 9

to 20°C, 9 to 15°C, 9 to 10°C, 10 to 15°C, typically 0 to 15°C, more typically 0 to 10°C, even more typically 3°C to 10°C, most typically 5°C to 10°C and even most typically 5°C. Alternatively the crushed or powdered metal carbonate, or metal carbonate hydroxide or metal oxide or mixture thereof is added to an aqueous solution of the 5 carbonic acid or hydrated carbon dioxide or carbon dioxide gas.




Typically one litre of water is placed in a container and sufficient carbonic acid or carbonated water or hydrated carbon dioxide or carbon dioxide gas is added to produce a pH value of approximately pH 5.2. (In practice, approximately 40 to 45mL of chilled (5°C) carbonated mineral water is used depending on the initial pH of the water). The 10 container is sealed and the contents are mixed. 485mg magnesium carbonate hydroxide pentahydrate powder $(\text{MgCO}_3)_4 \cdot \text{Mg}(\text{OH})_2 \cdot 5\text{H}_2\text{O}$, molecular weight 485 is added. The container is again sealed and the contents are mixed. The container is stored at a temperature of 0 to 10°C and the contents mixed regularly. Sufficient time is allowed for a clear solution of magnesium bicarbonate to develop at a range of pH 8.0 to pH 8.5, 15 typically pH 8.3. This takes approximately 24 to 72 hours. Alternatively the carbonic acid or carbonated water or hydrated carbon dioxide or carbon dioxide gas is added to the magnesium carbonate hydroxide pentahydrate powder in water.

Alternatively one litre of water is placed in a container and sufficient carbonic acid or carbonated water or hydrated carbon dioxide gas is added to produce a pH value less than 20 pH 5.2. (In practice, approximately 30mL to 40mL of chilled water is used depending on the initial pH of the water). The container is sealed and the contents are mixed. 485mg magnesium carbonate hydroxide pentahydrate powder $(\text{MgCO}_3)_4 \cdot \text{Mg}(\text{OH})_2 \cdot 5\text{H}_2\text{O}$, molecular weight 485 is added. The container is again sealed and the contents are mixed. The container is stored at a temperature of 0 to 10°C and the contents mixed regularly. 25 The pH of the water is then adjusted with an alkali such as sodium hydroxide or potassium hydroxide to a pH of 8 to 8.5, typically pH 8.3. Alternatively the carbonic acid or carbonated water or hydrated carbon dioxide or carbon dioxide gas is added to the magnesium carbonate hydroxide pentahydrate powder in water.


Generally once the solution is prepared, the solution may be stored in a closed container 30 under a blanket of carbon dioxide gas or a mixture of carbon dioxide gas and usually a nondeleterious inert gas, for example, argon, helium and/or nitrogen to maintain the solution at a pH of 7 to 9 and at a temperature of 0 to 25°C and at 0.8 to 5 atm. The

carbon dioxide gas blanket prevents loss of carbon dioxide from the solution. The amount of carbon dioxide in the gaseous mixture provides partial pressure on the liquid which is substantially equal to the partial pressure of carbon dioxide from carbon dioxide from the solution which is produced from equilibrium of bicarbonate in the solution at the particular temperature. In this way the solution is stabilised. If the solution were left in an open container for any substantial length of time precipitation of metal carbonate from the solution would occur as a result of decomposition of the bicarbonate in the solution as carbon dioxide is liberated from the solution. By using a stabilising agent in and/or above the solution such decomposition is substantially minimised or prevented. Alternatively the solution may be stored in a closed or sealed container (generally airtight) which is substantially filled with the solution whereby there is substantially no gas in the container or little gas compared to the amount of liquid in the container.

The relevant chemical reactions may be represented by the following equations:




and




pH 7.0 to pH 8.0



pH 8.0 to pH 8.3



pH 8.3 to pH 8.5



pH 8.5 to pH 9.0

Usually one litre of the magnesium bicarbonate solution prepared above contains approximately 120mg of magnesium per litre of aqueous neutral to mildly alkaline metal bicarbonate solution and approximately 600mg of bicarbonate. 500mg sodium bicarbonate (or potassium bicarbonate) is added to the magnesium bicarbonate solution and mixed. The mixture is stored in a sealed container in a refrigerator. The mixture

contains approximately 120mg magnesium per litre of aqueous neutral to mildly alkaline metal bicarbonate solution, 135mg sodium per litre of aqueous neutral to mildly alkaline metal bicarbonate solution and 950mg bicarbonate per litre of aqueous neutral to mildly alkaline metal bicarbonate solution.

- 5 In the body, normal intracellular pH value is pH 7.2. Under acidic conditions, such as adenosine triphosphate (ATP) hydrolysis, intracellular pH value may decrease to pH 6.5. In practice, a pH value is chosen for bicarbonate solutions that exceeds normal blood plasma pH value (pH > 7.38).

A low temperature, between 0 and 10°C, typically 5 to 10°C, ensures that carbon dioxide
10 stays dissolved in solution to maximise carbon dioxide hydration. Above 15 to 20°C, the solubility of carbon dioxide is low, the carbon dioxide leaves the solution, and particles and sediments may occur in the solution. Above 15 to 20°C, the solution may be cloudy in appearance.

At high magnesium concentrations, a minimum time, at least 24 to 72 hours at 5°C, is
15 required for completion of the kinetic processes that produce a clear solution of magnesium bicarbonate. (The kinetic processes include the hydration of carbon dioxide, the dissolution of magnesium carbonate and the dissolution of magnesium hydroxide.)

The concentration of magnesium cations (in association with bicarbonate anions) is generally in the range 25mg to 250mg per litre aqueous neutral to mildly alkaline metal
20 bicarbonate solution (depending on the pH value of the metal bicarbonate solution).

Usually the maximum magnesium concentration that can be maintained in solution as magnesium bicarbonate may be approximately 120mg per litre aqueous neutral to mildly alkaline metal bicarbonate solution at pH 8.3. As the pH value decreases, the concentration of magnesium that can be maintained in solution increases. Because
25 magnesium chloride is soluble, higher concentrations of magnesium can be maintained in solution if chlorides (such as sodium chloride) are added to the aqueous neutral to mildly alkaline metal bicarbonate solution.


The solubility product constant for magnesium carbonate is reported to be approximately 3.5×10^{-8} . The solubility product constant for magnesium hydroxide is reported to be
30 approximately 1.1×10^{-11} . Calculated from these values, the maximum concentrations of magnesium cations that can exist in solution as carbonates or hydroxides are

approximately 20mg per litre aqueous metal bicarbonate solution and 10mg per litre aqueous neutral to mildly alkaline metal bicarbonate solution respectively.


Generally the aqueous neutral to mildly alkaline metal bicarbonate solution of the invention is administered or consumed orally. Typically the solution is an orally drinkable solution. Typically the solution is a therapeutic orally drinkable solution. For example, the aqueous neutral to mildly alkaline metal bicarbonate solution may be prepared as a solution or an iced confectionary, such as an ice block or iced dessert, which is ingested orally. Alternatively the aqueous neutral to mildly alkaline metal bicarbonate solution may be prepared in the form of a tablet, lozenge or lolly which is ingested orally. For example, the aqueous neutral to mildly alkaline metal bicarbonate solution may be administered for metabolic acidosis or renal failure. Optionally the solution may be sterilised. Typically the aqueous neutral to mildly alkaline metal bicarbonate solution is prepared as a solution which is ingested hourly, daily, monthly or yearly. The amount of aqueous neutral to mildly alkaline metal bicarbonate solution administered in a day ranges from 250mL to 6 litres, typically 250mL to 5.5 litres, 250mL to 5 litres, 250mL to 4.5 litres, 250mL to 4 litres, 250mL to 3.5 litres, 250mL to 3 litres, 500mL to 6 litres, 500mL to 5.5 litres, 500mL to 5 litres, 500mL to 4.5 litres, 500mL to 4 litres, 500mL to 3.5 litres, 500mL to 3 litres, more typically 1 litre to 6 litres, 1 litre to 5.5 litres, 1 litre to 5 litres, 1 litre to 4.5 litres, 1 litre to 4 litres, 1 litre to 3.5 litres, even more typically 1 litre to 3 litres, 1.5 litres to 6 litres, 1.5 litres to 5.5 litres, 1.5 litres to 5 litres, 1.5 litres to 4.5 litres, 1.5 litres to 4 litres, 1.5 litres to 3.5 litres, most typically 2 to 3 litres, usually 2.1 to 3 litres. The aqueous neutral to mildly alkaline metal bicarbonate solution may be administered on a full or empty stomach, typically the aqueous neutral to mildly alkaline metal bicarbonate solution is administered on an empty stomach. Usually 1.5 to 3 litres, more typically 1.5 to 2.4 litres, even more typically 1.8 to 2.1 litres and usually between 1.8 and 2.7 litres of aqueous neutral to mildly alkaline metal bicarbonate solution is ingested on an empty stomach in approximately 300mL volumes at set times each day. The solution may be administered according to these latter dosages over short (for example 1 to 10 days) or long (for example 6 months to 10 years or more) periods as required. Usually the amount of aqueous neutral to mildly alkaline metal bicarbonate solution administered to a mammal is 5 to 100mL per Kg, more usual 10 to 50mL per Kg, most usual 14 to 29mL per Kg or 25 to 43mL per Kg.

Typically the aqueous neutral to mildly alkaline metal bicarbonate solution is consumed on an empty stomach. Usually consumption in this manner avoids the mixing of bicarbonate anions with stomach acid which may result in the loss of bicarbonate. Usually the aqueous neutral to mildly alkaline metal bicarbonate solution is consumed in small amounts at set times each day to avoid a rapid increase in the bicarbonate concentration of body fluids. Usually the amount of aqueous neutral to mildly alkaline metal bicarbonate solution consumed at commencement is 500mL per day and is increased by increments over a period of one month to the maximum consumption. This start-up schedule generally avoids any gastrointestinal side effects due to the smooth muscle relaxation properties of magnesium.

The advantages of the aqueous neutral to mildly alkaline metal bicarbonate solution of the invention are that the magnesium cations function as bicarbonate transporters into body cells. Magnesium bicarbonate enters body cells and the bicarbonate anions function to displace from equilibrium the dissociation reaction of intracellular carbonic acid. Magnesium bicarbonate enters body cells and the bicarbonate anions function as an intracellular proton sink (or proton scavenger). These reactions can be represented by the one equation



Magnesium bicarbonate enters body cells and the bicarbonate anions function to displace from equilibrium the hydration reaction of carbon dioxide which is catalysed by the enzyme carbonic anhydrase. This reaction can be represented by the equation




Usually appropriate salts of magnesium, sodium, potassium, calcium and lithium should not exceed the concentrations of the component elements recommended by health authorities. The concentrations of component elements cannot exceed concentrations restricted by the solubility product constants of respective hydroxides and carbonates.

Example 1

An experiment to decrease intracellular proton concentrations and to increase intracellular bicarbonate concentrations in mammalian cells *in vitro*

Aqueous bicarbonate anions act as proton sinks in the presence of excess proton concentrations in solution. This reaction can be represented by the chemical equation



5 In the presence of sufficient concentrations of bicarbonate anions, the reaction is essentially complete and proton concentrations decrease. The pH value of the solution increases. When plasma bicarbonate anions are present outside mammalian body cells in sufficient concentrations, they are translocated into the cytoplasm of the cells across the cell plasma membranes. Indeed, bicarbonate anions equilibrate rapidly across mammalian
10 cell membranes. Bicarbonate translocation into cells takes place via several processes. These processes include a chloride-bicarbonate anion exchange and a sodium dependent chloride-bicarbonate anion exchange and potassium co-transport and magnesium co-transport.

An experiment was conducted to decrease intracellular proton concentrations and to
15 increase intracellular bicarbonate concentrations in mammalian body cells *in vitro*. Throughout the experiment, extracellular pH determinations were made using a pH electrode and intracellular pH determinations were made using a trapped fluorescein derivative. An increase in intracellular proton concentrations (intracellular acidification) was achieved by applying 10 mmol ammonium chloride (NH₄Cl) solution to a suspension
20 of cells and then removing the NH₄Cl. An increase in intracellular bicarbonate concentrations was achieved by applying an aqueous metal bicarbonate solution to a suspension of cells. The aqueous metal bicarbonate solution contained approximately Mg²⁺ 120 mg per litre, Na⁺ 135 mg per litre and HCO₃⁻ 950 mg per litre at pH 8.3. This aqueous metal bicarbonate solution was equivalent to 15 mmol bicarbonate
25 approximately. Blood was collected in sodium heparin from a range of mammals and the leucocytes removed. The leucocytes were washed and re-suspended in isotonic saline.

Intracellular pH determinations were made by loading leucocytes for 15 minutes with (10 micromol in saline) 2,7-bis(carboxyethyl)-5,6-carboxyfluorescein (BCECF). Cells were illuminated at 440 nm and 490 nm and fluorescence was measured at 530 nm.

30 The experiment utilising sheep leucocytes is given stepwise below:

Step 1. Increase in intracellular proton concentrations (cytoplasmic acidification)

A. Leucocytes suspended in normal saline after pretreatment with fluorescein.

Extracellular pH 7.2

Intracellular pH 7.1

5 B. 10 mmol ammonium chloride (NH_4Cl) solution pH 7.5 applied to suspension of leucocytes for 10 minutes.

C. Leucocytes washed and re-suspended in normal saline.

Extracellular pH 7.3

Intracellular pH 6.1

10 **Result:** Cells have increased intracellular proton concentrations. Cytoplasm is acidified.

Step 2. Decrease in intracellular proton concentrations

A. Acidified leucocytes (from Step 1.) divided into two groups; Control group and Treatment group.

15 B. Treatment group of leucocytes exposed to aqueous metal bicarbonate solution. After 3 minutes:

Extracellular pH 7.5

Intracellular pH 7.0

C. Control group of leucocytes not exposed to aqueous metal bicarbonate solution.

20 After 5 minutes:

Extracellular pH 7.2

Intracellular pH 6.6

25 **Result:** Cells treated with aqueous metal bicarbonate solution rapidly decrease intracellular proton concentrations. Cytoplasm shows rapid recovery from acidification relative to non-treated cells.

Step 3. Increase in intracellular bicarbonate concentrations

A. Leucocytes suspended in normal saline after pretreatment with fluorescein.

Extracellular pH 7.2

Intracellular pH 7.1

- B. Aqueous metal bicarbonate solution applied to suspension of leucocytes for 20 minutes.

Extracellular pH 7.9

Intracellular pH 7.4

- 5 **Result:** Cells treated with aqueous metal bicarbonate solution have increased intracellular bicarbonate concentrations which are manifested by an increase in pH value of cytoplasm.

The experiment was repeated with leucocytes from mice, rats, guinea pigs, cattle, horses, dogs, cats and humans. In all cases, acidified cells treated with aqueous metal bicarbonate solution had decreased intracellular proton concentrations. In all cases, cells treated with aqueous metal bicarbonate solution had increased intracellular bicarbonate concentrations which were manifested by increased pH values of cytoplasm.

The experiment was repeated with aqueous metal bicarbonate solutions that contained a range of concentrations of Mg^{2+} , Na^+ , HCO_3^- and K^+ and Ca^{2+} ions. Significant results were obtained for the following range of concentrations:

Ion	Range of concentrations to achieve significant results		
Mg^{2+}	20	to	120 mg/litre
Na^+	50	to	500 mg/litre
K^+	50	to	500 mg/litre
Ca^{2+}	20	to	150 mg/litre
HCO_3^-	250	to	2,100 mg/litre
(HCO_3^-)	(4 mmol to 35 mmol)		

Significant results were obtained for pH range pH 7.5 to 9.5. (pH 9.5 was achieved by the addition of NaOH).

Aqueous metal bicarbonate solutions, containing a range of cation and bicarbonate anion concentrations, decrease intracellular proton concentrations and increase intracellular bicarbonate concentrations in mammalian cells *in vitro*.

Example 2

An experiment to demonstrate bicarbonate anion translocation from aqueous metal bicarbonate solution into the mammalian body against a bicarbonate anion concentration gradient

Mammalian plasma contains bicarbonate anions at a concentration about 25 mmol (HCO_3^- 1,500 mg per litre). When ingested, aqueous metal bicarbonate solution produces biochemical, physiological and medical effects at bicarbonate anion concentrations about 16 mmol (HCO_3^- 950 mg per litre). Aqueous metal bicarbonate solution, at bicarbonate anion concentration about 16 mmol, contains two thirds the bicarbonate anion concentration of plasma, so bicarbonate anions must be translocated into the mammalian body against a bicarbonate anion concentration gradient.

Mammalian plasma contains cations at concentrations around Mg^{2+} 24 mg per litre, Na^+ 3,300 mg per litre, K^+ 175 mg per litre and Ca^{2+} 100 mg per litre. Aqueous metal bicarbonate solution commonly contains cations at concentrations around Mg^{2+} 120 mg per litre, Na^+ 135 mg per litre, K^+ 100 mg per litre and Ca^{2+} 20 mg per litre. Aqueous metal bicarbonate solution commonly contains 5 times the magnesium cation concentration of plasma. Other cations are present commonly in aqueous metal bicarbonate solution in concentrations lower than plasma.

The concentrations of cations and anions in plasma can be compared with concentrations of cations and anions in aqueous metal bicarbonate solution by examination of the following table:

Concentrations of cations and anions		
Ion	Plasma	Aqueous metal bicarbonate solution
Cl^-	3,600 mg/litre	0 mg/litre
Na^+	3,300 mg/litre	135 mg/litre
HCO_3^-	1,500 mg/litre	950 mg/litre
K^+	175 mg/litre	100 mg/litre
Ca^{2+}	100 mg/litre	20 mg/litre
Mg^{2+}	24 mg/litre	120 mg/litre

It is suggested that sodium cations and chloride anions leave plasma along their respective concentration gradients and magnesium and bicarbonate ions enter plasma along a

magnesium cation concentration gradient. Magnesium functions as a bicarbonate transporter. In addition, it is suggested that bicarbonate anions enter plasma by chloride-bicarbonate exchange processes along a chloride anion concentration gradient (chloride 'out', bicarbonate 'in').

5 In mammals, any large increases in plasma bicarbonate concentrations can be decreased normally by a number of biochemical and physiological homeostatic control processes. These processes occur in time frames that range from minutes to hours and longer. One of the main control processes that occurs as a result of increased plasma bicarbonate concentration is an alteration in bicarbonate chemistry in the kidneys. This is manifested
10 by a decrease in proton concentration in urine and by a pH value of urine that is less acidic. In the presence of increased plasma bicarbonate, kidney tubule cells decrease their excretion of protons. Kidney control of bicarbonate concentration is not instantaneous and occurs within a time frame of several hours to several days. Unless a mammal has physiological or clinical acidosis, it is difficult to detect small increases in plasma
15 bicarbonate concentration. Any increases in plasma bicarbonate concentration are taken up by body cells. Indeed, plasma bicarbonate equilibrates with intracellular bicarbonate rapidly. In a normal mammal, a measurable increase in plasma bicarbonate concentration occurs only during an artificially induced alkalosis and is detectable either when the consumption of bicarbonate anions (as NaHCO_3) greatly exceeds the concentration of
20 bicarbonate in normal plasma or when bicarbonate anions (as NaHCO_3) are administered intravenously.

An experiment was conducted to determine if bicarbonate anions in aqueous metal bicarbonate solutions are translocated against a bicarbonate concentration gradient into the body. Bicarbonate translocation against a concentration gradient could occur either via
25 energy (ATP) dependent processes or via anion (chloride-bicarbonate) exchange or via co-transport with cations along cation concentration gradients. There are also complex thermodynamic processes involving intracellular and extracellular concentrations of bicarbonate anions, hydroxide anions, protons and carbon dioxide that may assist in the overall translocation of bicarbonate anions. These processes often involve the production
30 of bicarbonate anions by carbonic anhydrase enzymes. In the experiment, entry of bicarbonate anions into the body was assessed by determinations of proton concentration in urine; that is, the pH value of urine.

Ten people had urine pH value assessed once per week for 3 months. Urine pH values were assessed once per week for a further 3 months after commencement of consumption of aqueous metal bicarbonate solution. The aqueous metal bicarbonate solution contained approximately Mg^{2+} 120 mg per litre, Na^+ 135 mg per litre and HCO_3^- 950 mg per litre.

5 The major component of the solution was magnesium bicarbonate $\text{Mg}(\text{HCO}_3)_2$ 720 mg per litre approximately. Results are given below:

Mean pH value of urine

(Early morning sample)

Prior to consumption of aqueous

10 metal bicarbonate solution:

pH 5.9

After commencement of consumption

of aqueous metal bicarbonate solution:

pH 6.7

The consumption of aqueous metal bicarbonate solution decreases proton excretion by the kidneys. The pH value of urine increases.

15 These results demonstrate that bicarbonate anions from aqueous metal bicarbonate solution are translocated against a bicarbonate anion concentration gradient into the body. This may occur either via co-transport with cations along a cation concentration gradient or via chloride-bicarbonate exchange processes along a chloride anion concentration gradient (chloride 'out', bicarbonate 'in'). In the case of aqueous metal bicarbonate solution, the

20 only cation concentration gradient possible is that involving magnesium cation concentrations.

The consumption of aqueous metal bicarbonate solution leads to an increase in bicarbonate anion concentration in the body which is manifested by a decrease in proton concentration in urine; an increase in pH value of urine.

25


Example 3

An experiment to improve the buffering capacities of the extracellular and intracellular bicarbonate buffers and to decrease senescence and to increase longevity in a representative mammal

Mammalian body cells produce continuously concentrations of carbon dioxide. Upon

30 hydration, carbon dioxide increases proton concentrations in the cytoplasm of body cells. The pH values of the cytoplasm of body cells are lowered. The production of protons in

cytoplasm by the hydration of carbon dioxide can be represented by the following chemical equations:



5 The protons produced in the cytoplasm of body cells by the hydration of carbon dioxide, and other intracellular reactions, are buffered normally by intracellular bicarbonate buffers. The bicarbonate anions in intracellular buffers derive mainly from the extracellular bicarbonate of blood plasma. The bicarbonate anions in blood plasma originate from erythrocytes as products of erythrocyte carbonic anhydrase enzyme
10 reactions.

When plasma bicarbonate anions are present outside mammalian body cells in sufficient concentrations, they are translocated into the cytoplasm of the cells across the cell plasma membranes. Indeed, plasma bicarbonate equilibrates with cytoplasmic bicarbonate rapidly. Bicarbonate translocation into cells takes place via several processes. These
15 processes include a chloride-bicarbonate anion exchange and a sodium dependent chloride-bicarbonate anion exchange and potassium co-transport and magnesium co-transport. There are also complex thermodynamic processes involving intracellular and extracellular concentrations of bicarbonate anions, hydroxide ions, protons and carbon dioxide that may assist in the overall translocation of bicarbonate anions. These processes often involve the
20 production of bicarbonate anions by carbonic anhydrase enzymes.

Concentrations of bicarbonate anions that are translocated into mammalian body cells improve the buffering capacity of the cytoplasm of the cells. Concentrations of bicarbonate anions and concentrations of carbon dioxide form a buffer system described by the Henderson-Hasselbalch equation:

$$25 \quad \text{pH} = \text{pK} + \log ([\text{HCO}_3^-]/[\text{H}_2\text{CO}_3])$$


(Where pK is the pK of hydrated carbon dioxide H_2CO_3 and has an approximate numerical value of 6.35)

For a classical (closed system) buffer to be effective, the ratio of the conjugate base to the acid (in the above case $[\text{HCO}_3^-]/[\text{H}_2\text{CO}_3]$) must be between 0.1 and 10. This ratio applies
30 also to buffers in biological (open) systems. In mammalian body cells, the continuous and open production of carbon dioxide means that continuous supplies of bicarbonate anions

are required to maintain effective and optimal buffering capacities. Under conditions of excess proton concentrations, from carbon dioxide production and ATP hydrolysis and other metabolic processes, the supply of bicarbonate fails and the effective and optimal buffering capacities of mammalian body cells falter.


- 5 The vitality of mammalian body cells is linked critically to the buffering capacities of the extracellular fluids and the cytoplasm of the cells. Processes of cellular degeneration occur when buffering capacities falter in the presence of excess proton concentrations. Cellular degenerations are manifested in the mammalian body by degenerative diseases and senescence. Examples of degenerative diseases in mammals that are linked casually
- 10 to extracellular and intracellular proton concentrations include osteoporosis, osteoarthritis, the diseases associated with chronic inflammation, the diseases associated with lysosomal enzyme activities, the diseases associated with oxidations of cell nucleic acids, cell protein amino acids and cell membrane lipids, and the diseases associated with aberrations of mitochondrial respiration.
- 15 An experiment was conducted to improve the buffering capacities of the extracellular and intracellular bicarbonate buffers and to consequently decrease senescence and increase longevity in a representative mammal. One hundred and ten Merino ewe lambs were divided randomly at weaning into a control group and a treatment group. The groups were of equal size and were maintained under similar conditions except for the pH values
- 20 and aqueous metal bicarbonate concentrations of drinking water supplies. Sheep were selected as the representative mammal because their life span and body weight are more representative of typical mammals than laboratory rodents, their life span is not excessively long, their body size permits multiple blood and tissue sample collections, they are easy to handle and their husbandry is suited to experimental conditions. The
- 25 control group was maintained, for the full life span of the sheep, in small experimental paddocks with slightly acidic (less than pH 6.5) drinking water supplies that contained bicarbonate concentrations less than 30 mg per litre. The treatment group was maintained, for the full life span of the sheep, in small experimental paddocks with slightly alkaline (pH 7.8 to 9.0) drinking water supplies that contained bicarbonate concentrations between
- 30 300 mg per litre and 800 mg per litre. The drinking water supplies for the treatment group were loaded with the appropriate concentrations of bicarbonate anions by the addition of crushed and powdered magnesite MgCO_3 to the water. The magnesite frequently contained calcite CaCO_3 and dolomite $(\text{Ca,Mg})\text{CO}_3$. The magnesite was dissolved in the

drinking water either with the assistance of commercial supplies of carbon dioxide gas or carbonic acid or with local supplies of hydrated carbon dioxide. This dissolution process can be represented by the following chemical equations:




The treatment group of sheep consumed slightly alkaline (pH 7.8 to 9.0) drinking water that contained bicarbonate concentrations between 300 mg per litre and 800 mg per litre.

At this pH value, and this bicarbonate concentration, bicarbonate was mostly in the form of magnesium bicarbonate $\text{Mg}(\text{HCO}_3)_2$:



In addition, some sediments of carbonate $(\text{Ca},\text{Mg})\text{CO}_3$ were present in the drinking water during summer months:



The mean pH values and the mean magnesium, calcium and bicarbonate concentrations in the drinking water supplies are given below (the concentrations of cations and bicarbonate anions were not stoichiometric in the drinking water - particularly the drinking water of the control group - because of the presence of some concentrations of sulphate, chloride and sodium ions):

Means of parameters in drinking water

	Control Group	Treatment Group
pH	6.1	8.4
Mg^{2+} mg/litre	13	110
Ca^{2+} mg/litre	20	30
HCO_3^- mg/litre	25	660

In the late stages of pregnancy, there is a tendency for pregnant mammals to become hypoglycaemic and hyperketonaemic. Hyperketonaemia subjects the pregnant mammal to an acid load (increase in proton concentrations). This acid load may result in clinical acidosis. Like all mammals, pregnant ewes tend to be hypoglycaemic and hyperketonaemic late in pregnancy. In ewes affected clinically with acidosis, bicarbonate concentrations range between 14 to 20 mmol per litre plasma.

Over several years, plasma bicarbonate concentrations were determined for the control group and the treatment group one week prior to lambing. Determination of plasma bicarbonate concentrations prior to lambing is a direct measure of extracellular and intracellular bicarbonate buffering capacity. In ewes with effective extracellular and intracellular bicarbonate buffers, bicarbonate concentrations are maintained in a range between 24 to 27 mmol per litre plasma. Plasma bicarbonate concentrations are given below:

Mean plasma bicarbonate concentrations one week prior to lambing (mmol per litre)

Age (years)	Control Group	Treatment Group
4	24.9	26.1
6	22.8	25.9
8	22.2	26.4
10	21.9	25.8

The treatment group had larger plasma bicarbonate concentrations than the control group.

The consumption of aqueous metal bicarbonate solution, principally magnesium bicarbonate solution, improves the buffering capacities of extracellular and intracellular bicarbonate buffers in mammals.

In mammalian demography, there are two measurements utilised commonly in the experimental study of degenerative diseases and senescence. The first measurement is called fifty percent survival. Fifty percent survival describes the chronological age at which half an original population has died. The second measurement is called maximum life span. Maximum life span describes the age of the longest lived survivors of a population. The fifty percent survival measurement is considered to reflect susceptibility to accidents and infectious and degenerative diseases in mammals. The maximum life span measurement is considered to reflect the innate processes of senescence in

mammals. The fifty percent survival measurement and the maximum life span measurement for the control group and the treatment group are given below:

Fifty percent survival

Control group 8 years

Treatment group 11 years

Maximum life span

Control group 13 years

Treatment group 17 years

The treatment group had a larger fifty percent survival measurement and a larger maximum life span measurement than the control group.

The death of each member of a population of mammals can be plotted graphically. The continuous function representing mortality in a population is known as a survival curve. Survival curves for the control group and treatment group are represented in Figure 1.

The survival curves show that more mature sheep were alive in the treatment group than the control group at any time. This occurred with the consumption of normal physiological volumes of water (as aqueous metal bicarbonate solution).

The consumption of aqueous metal bicarbonate solution, extends the maximum life span of mammals by at least twenty percent and increases the number of mature mammals alive at any time.

Senescence in mammals is characterised by progressive oxidations of the structural and function molecules that constitute body cells and tissues. These oxidations occur particularly in nucleic acids, protein amino acids and cell membrane lipids.

Because protons often participate in biological redox reactions, oxidations of many structural and functional molecules in body cells and tissues are increased in rate by the presence of excess proton concentrations. Oxidations of structural and functional molecules are increased in rate by acidic conditions.

In general, oxidations of molecules are linked to proton concentrations described by a formulation of the Gibbs energy equation

$$E_{pH} = E_m + 2.3 RT/F \log ([\text{oxidised}]/[\text{reduced}])$$

where E_{pH} is a measure of oxidising power at a particular pH value and E_m is the mid-point potential. In practice, E_{pH} is decreased by between -30 mV and -60 mV for each decrease in proton concentration by a factor of 10. That is, oxidising power is decreased by between -30 mV and -60 mV for each increase in pH value by 1 pH unit.

- 5 Oxidations of nucleic acids and protein amino acids lead to nucleic acid and protein degradation respectively. These degradations lead to senescence in mammals. Nucleic acid degradation is manifested by either cell death or cell transformation to the cancerous state. Protein degradation is manifested by increased urea concentrations in the body which can be detected in the plasma.
- 10 Determination of plasma urea concentrations in elderly mammals is a direct measure of amino acid oxidation, protein degradation and overall nitrogen (anabolic/catabolic) balance. Determination of plasma urea concentrations in elderly mammals is a direct measure of cellular degenerations and senescence.

Over several years, plasma urea concentrations were determined for the control group and
15 the treatment group. Plasma urea concentrations are given below:

Mean plasma urea concentrations in elderly sheep (mmol per litre)

Age (years)	Control Group	Treatment Group
8	11	5
10	13	3
12	13	7

The treatment group had smaller plasma urea concentrations than the control group.

- The consumption of aqueous metal bicarbonate solution, principally magnesium bicarbonate solution, decreases amino acid oxidations, decreases protein degradation
20 and improves overall nitrogen (anabolic/catabolic) balance in mammals. The consumption of aqueous metal bicarbonate solution, principally magnesium bicarbonate solution, delays cellular degenerations and senescence in mammals.

Autopsies were performed on sheep, when conditions permitted, within 24 hours of death.

Macroscopic signs of significant degenerative diseases and other diseases were recorded.

- 25 Significant pathology is given below:

Prevalence of pathology at autopsy (%)

Macroscopic Significant Pathology	Control Group	Treatment Group
(* most significant)	(42 autopsies)	(38 autopsies)
Lungs	24%	21%
*Heart	29%	11%
Liver	43%	21%
Kidney	24%	16%
Other Genito-urinary	17%	16%
Lymph nodes	40%	37%
Intestinal tract	10%	8%
*Joints	43%	5%
*Bone	24%	3%
Teeth	71%	40%
*Skin-wool	48%	21%
Cancer	12%	3%

The treatment group had a lower overall prevalence of pathology than the control group. In general, pathology in the treatment group was delayed (sheep were older at autopsy) and progression was less advanced.

The consumption of aqueous metal bicarbonate solution, principally magnesium bicarbonate solution, decreases the prevalence of joint pathology (arthritis) and bone pathology (osteoporosis) and cardiac pathology and skin pathology most significantly and decreases the overall prevalence of the pathology of most organs.

Example 4

10 An experiment to distinguish between the consumption of magnesium bicarbonate and the consumption of magnesium cations *per se* in increasing longevity in a mammal

15 An experiment was conducted to assess if the consumption of magnesium bicarbonate increased longevity in a mammal compared to the consumption of magnesium cations *per se*. A short-lived mammalian species was chosen. Short-lived mammals possess high levels of proton leak across inner mitochondrial membranes, high levels of carbonic

anhydrase enzyme activities (for acid production) and high levels of spontaneous cancer development and spontaneous death. Any increase in longevity in a short-lived species is indicative of an improvement in fundamental cell biochemistry. Two hundred outbred (Swiss) female mice were divided randomly at weaning into two groups of 100 mice and 5 were maintained under identical management and environmental conditions.

One group of mice was supplied with drinking water that consisted of aqueous metal bicarbonate solution with a pH value between pH 8.1 and pH 8.5. The aqueous metal bicarbonate solution contained approximately Mg^{2+} 120mg per litre, Na^+ 135mg per litre and HCO_3^- 950mg per litre. The major component of the solution was magnesium bicarbonate $\text{Mg}(\text{HCO}_3)_2$ 720mg per litre approximately. The second group of mice was supplied with drinking water that contained magnesium sulphate (Epsom salts) 1 gram per litre with a pH value between pH 6.5 and pH 7.0. This drinking water contained approximately Mg^{2+} 120mg per litre. Bicarbonate anions were absent.

Both groups of mice were fed commercial laboratory food that contained 1 gram of magnesium per kilogram of food. Both groups of mice were fed on alternate days with no food available on the other days. Group-specific drinking water (as described above) was available at all times. Feeding on alternate days decreased the possible loss of bicarbonate anions by stomach acid and food ingesta.

Results of the experiment are given below:

20

Fifty Percent Survival

Group consuming magnesium bicarbonate	790 days
Group consuming magnesium sulphate	736 days

Maximum Life Span

Group consuming magnesium bicarbonate	1152 days
Group consuming magnesium sulphate	1040 days

The group of mice consuming aqueous metal bicarbonate solution had longevity increased by ten percent compared to the group of mice consuming magnesium cations *per se*.


The consumption of aqueous metal bicarbonate solution, principally magnesium bicarbonate solution, extends the maximum life span of mammals by ten percent more than the consumption of magnesium cations *per se*.

Example 5

An experiment to decrease the clinical signs of osteoarthritis

Osteoarthritis is a disease of degeneration. There is degradation and inflammation of the joints of the body. Osteoarthritis is defined as a disease process involving a disturbance of the normal balance of degradation and repair in the articular cartilage and subchondral bone of joints. This disturbance of balance causes areas of morphological damage and results in clinical problems such as pain and disability. Osteoarthritis is manifested as a slowly progressive degeneration of the joints of the hands and large weight-bearing joints (hips and knees). It is common in post menopausal women. Osteoarthritis is characterised by pain, enlargement of joints and limitation of joint movements. The linings of osteoarthritic joints show a moderate to marked degree of inflammation. The principle pathological changes associated with osteoarthritis are destruction of joint cartilage and neoformations of bone at joint margins (osteophytes). In osteoarthritis, destruction of joint cartilage is caused by acid protease enzymes (and other enzymes) derived often from the lysosomes of cartilage cells (chondrocytes), inflammatory cells and other cells.

Acid protease enzymes possess optimal activity in an acidic environment; that is, an environment with high proton concentrations. Proton concentrations involved in the pathogenesis of osteoarthritis derive from the hydration of carbon dioxide catalysed by intracellular carbonic anhydrase enzymes. The production of protons by carbonic anhydrase enzymes can be represented by the equation



Protons formed by carbonic anhydrase enzymes are concentrated by intracellular V-type proton pumps and stored in the endosomes and lysosomes of body cells.

Functional endosomes and lysosomes maintain internal concentrations of protons which give them internal pH values between pH 3.0 and pH 6.0. Many degenerative diseases, including osteoarthritis, involve intracellular and extracellular release of lysosomal enzymes. In osteoarthritis, chemical fluxes through the reactions catalysed by lysosomal enzymes result in the breakdown of cartilage and bone.

An experiment was conducted to assess if the clinical signs of osteoarthritis could be decreased by the consumption of aqueous metal bicarbonate solution. The clinical signs of osteoarthritis include pain, swelling, inflammation, skin discoloration, joint deformities

and decrease in joint function. An increase in extracellular and intracellular bicarbonate anion concentrations would decrease the production of protons from reactions catalysed by carbonic anhydrase enzymes, decrease the pumping of protons by V-type proton pumps, decrease the activities of acid protease enzymes and decrease other activities of lysosomes.

5 The clinical signs of osteoarthritis would be alleviated. A group of ten people were chosen who had been diagnosed with having osteoarthritis. Each person in the group had been suffering from (clinical) osteoarthritis for between 2 and 5 years. Five of the group were post menopausal women who had clinical signs of osteoarthritis in the joints of their hands. The osteoarthritic joints included the distal and proximal interphalangeal joints of
10 the fingers and the carpometacarpal joint of the thumbs. In all 5 cases, loss of joint function was moderate to severe.

In all 5 cases, the women suffered pain, swelling of the fingers and loss of joint movement. Mucous cysts were associated with distal joint osteoarthritis. Lateral deformities occurred in some proximal joints with severe loss of joint function. Women
15 with affected thumbs had considerable loss of function and considerable pain. Many hands were "claw-like" in appearance (Fig. 2). The remainder of the group had osteoarthritis in the hips and knees. These people suffered pain and moderate loss of joint functions.

The people consumed aqueous metal bicarbonate solution with a pH value between pH 8.1
20 and 8.5. The aqueous metal bicarbonate solution contained approximately Mg^{2+} 120 mg per litre, Na^+ 135 mg per litre and HCO_3^- 950 mg per litre. The major component of the solution was magnesium bicarbonate $Mg(HCO_3)_2$ 720 mg per litre approximately.

Consumption of the aqueous metal bicarbonate solution was commenced at half a litre per day and increased by increments over a period of one month to between 2 to 3 litres per
25 day. Consumption occurred on an empty stomach to avoid the loss of bicarbonate by stomach acid (HCl). Consumption occurred in small amounts (300 ml) at set times each day to avoid rapid increases in bicarbonate concentrations of body fluids and to avoid over hydration.

The results of the experiment were unequivocal. Within 3 to 6 months, all participants in
30 the experiment demonstrated substantial decreases in the clinical signs of osteoarthritis.

In all cases, there were remissions in the clinical signs of osteoarthritis which were quantifiable by standard tests of movement, flexibility and strength. The participants

showed considerable increases in joint functions and decreases in acute and chronic joint swellings. The "stabbing" pain of osteoarthritis was alleviated. Some participants had remissions of inflammation and arthritis to the stage where many chronic swellings were no longer observable and joint mobilities and functions were restored (Fig. 3).

- 5 People in the experiment consumed aqueous metal bicarbonate solution continuously for at least 2 years. During this period, there was evidence of progressive improvement in healing processes. Mucous cysts associated with distal joint osteoarthritis were no longer visible.

Remissions in the clinical signs of osteoarthritis were maintained only with the continual
10 consumption of aqueous metal bicarbonate solution. Once the consumption of aqueous metal bicarbonate solution was halted, clinical signs of pain and swelling began to reappear within 10 days. Clinical signs again went into remission upon continuation of consumption of aqueous metal bicarbonate solution.

The consumption of aqueous metal bicarbonate solution, principally magnesium
15 bicarbonate solution, results in remissions in the clinical signs of osteoarthritis.


Example 6

An experiment to maintain and improve motor activity in mammals. An experiment to decrease fatigue and lethargy and improve motor activity. An experiment to decrease the fatigue and lethargy of chronic disease and improve motor activity

- 20 Mammals convert food energy into chemical energy that can be used by body cells to maintain essential cell processes and cell functions. The main chemical energy in mammalian body cells is the chemical ATP (adenosine triphosphate). ATP is synthesised mainly in the mitochondria of body cells. Mitochondrial ATP production is linked intimately to the respiration rates of mitochondria. The respiration rates of mitochondria
25 are dependent on many factors including the proton concentrations (pH values) of the cytoplasm of body cells. If the intracellular bicarbonate buffer of mammalian body cells is not maintained, and is not functional, proton concentrations increase in the cytoplasm and the pH value of the cytoplasm decreases. When proton concentrations increase in the cytoplasm sufficiently (pH value decreases sufficiently) the respiration rates of
30 mitochondria are diminished. When the respiration rates of mitochondria are diminished, the production of ATP is diminished. When the production of ATP is diminished, ATP concentrations in the cell decrease and the main chemical energy source for mammalian

body cells becomes depleted. Under these conditions, body cells cannot maintain essential cell processes and cell functions. The body becomes fatigued and lethargic.

In addition to the hydration of carbon dioxide *per se*, one of the sources of increased proton concentrations in the cytoplasm of body cells is the hydrolysis of ATP. The hydrolysis of ATP can be represented by the chemical equation



Increased proton concentrations from the hydrolysis of ATP occur particularly in the cytoplasm of muscle cells during muscular (motor) activity. This is referred to often as an increase in 'lactic acid' (the lactic acid is, in fact, lactate derived from glycolysis and the 'acid' is the protons derived from ATP hydrolysis).

An experiment was conducted to assess if motor activity could be maintained and improved in mammals by improving the buffering capacity of the extracellular and intracellular bicarbonate buffers.

Two hundred inbred (Balb c) female mice were divided randomly at weaning into two groups of 100 mice and maintained under identical conditions for 3 years. Control groups of mice were given drinking water that was deionised and slightly acidic (pH 5.0). Treatment groups of mice were given drinking water that consisted of aqueous metal bicarbonate solution with a pH value between pH 8.1 and 8.5. The aqueous metal bicarbonate solution contained approximately Mg^{2+} 120 mg per litre, Na^+ 135 mg per litre and HCO_3^- 950 mg per litre. The major component of the solution was magnesium bicarbonate $\text{Mg}(\text{HCO}_3)_2$ 720 mg per litre approximately.

Motor activity in mice was assessed at regular intervals for a 12 month period between 1 year and 2 years of age. Results of the experiment are given below:

Mean motor activity in mice

	Control Group	Treatment Group
Mean number of mice per hour climbing to lid of cage	26	95
Mean number of mice per hour engaged in exploratory activity	48	82

Mean time to exhaustion during enforced motor activity	5 minutes	9 minutes
--	-----------	-----------

The treatment group had improved motor activity relative to the control group.

The consumption of aqueous metal bicarbonate solution, principally magnesium bicarbonate solution, maintains and improves motor activity in mammals.

Mitochondria are described as 'efficient' if they maintain sufficient production of ATP for maintenance of essential cell processes and cell functions. Efficient mitochondria are the mitochondria of young mammals.

In mammals, there are declines in the efficiencies of mitochondria which are correlated to chronological age. The capacities of cells to maintain their particular energy requirements are diminished progressively with chronological age. Cells that are unable to meet their particular energy requirements undergo senescence, become non-functional and decline progressively towards cell death. This is manifested by body senescence and ageing.

Mitochondria are described as 'inefficient' if they cannot maintain the necessary production of ATP for maintenance of essential cell processes and cell functions. Mitochondrial inefficiency arises from oxidative damage to mitochondrial nucleic acids, mitochondrial enzymes and mitochondrial membrane proteins and lipids. Inefficient mitochondria gradually and progressively dominate in body cells through middle age to old age. Middle aged and elderly mammals are fatigued and lethargic relative to the young. Normal body cells attempt to produce buffers that maintain a cytoplasmic pH value of about pH 7.2. *In vitro*, if mitochondria are placed for a period in a medium either with an improved buffer at pH 7.2 or with a pH value buffered slightly higher than pH 7.2, there occurs an increase in mitochondrial respiration rate and an increase in the production of ATP.

An experiment was conducted to assess if fatigue and lethargy could be decreased and motor activity improved by improving the buffering capacity of the cytoplasmic bicarbonate buffer in a group of middle aged and elderly people.

Improving the buffering capacity of the cytoplasmic bicarbonate buffer would increase mitochondrial respiration rate and increase the production of ATP. Mitochondria would become more 'efficient'. More chemical energy would be available for maintenance of essential cell processes and cell functions. Fatigue and lethargy would decrease and motor activity would improve.

A group of nineteen people were chosen, with a mean age of 61 years, who had a history of fatigue and lethargy. In the context of this experiment, fatigue and lethargy were determined as subjective feelings of general exhaustion which were manifested by mild to moderate lack of function. The people consumed aqueous metal bicarbonate solution with a pH value between pH 8.1 and 8.5. The aqueous metal bicarbonate solution contained approximately Mg^{2+} 120 mg per litre, Na^{+} 135 mg per litre and HCO_3^{-} 950 mg per litre. The major component of the solution was magnesium bicarbonate $\text{Mg}(\text{HCO}_3)_2$ 720 mg per litre approximately.


Consumption of the aqueous metal bicarbonate solution was commenced at half a litre per day and increased by increments over a period of one month to between 2 to 3 litres per day. Consumption occurred on an empty stomach to avoid the loss of bicarbonate by stomach acid (HCl). Consumption occurred in small amounts (300 ml) at set times each day to avoid rapid increases in bicarbonate concentrations of body fluids and to avoid over hydration.

The results of the experiment were unequivocal. Within 3 months, all participants in the experiment demonstrated substantial decreases in fatigue and lethargy. All participants described a feeling of well-being (mild euphoria). All participants demonstrated an increased capacity for mild physical activity; an improvement in motor activity. Function was restored.

The consumption of aqueous metal bicarbonate solution, principally magnesium bicarbonate solution, decreases fatigue and lethargy and improves motor activity in middle aged and elderly people.

Chronic disease (including degenerative disease) is manifested often by chronic fatigue and lethargy and chronic pain. This is true particularly for chronic inflammatory diseases and autoimmune diseases.

The fatigue, lethargy and pain of chronic disease are correlated often to the high proton concentrations involved in the pathogenesis of chronic disease. In addition to the hydration of carbon dioxide *per se*, proton concentrations involved in the pathogenesis of chronic disease derive from the hydration of carbon dioxide catalysed by intracellular carbonic anhydrase enzymes. The production of protons by carbonic anhydrase enzymes can be represented by the equation



Protons formed by carbonic anhydrase enzymes are concentrated often by V-type proton pumps and stored in endosomes and lysosomes in the cell. The breakdown of endosomes and lysosomes creates concentrations of protons in the cell cytoplasm. This lowers the pH value of the cytoplasm and decreases the production of ATP in mitochondria. Cells become energy deficient. Cells are unable to maintain essential cell processes and cell functions. the body becomes fatigued and lethargic.

Functional endosomes and lysosomes maintain internal concentrations of protons which give them internal pH values between pH 3.0 and pH 6.0. Many of the chronic and degenerative diseases of the body involve intracellular lysosomal activities and intracellular and extracellular release of lysosomal enzymes. Chemical fluxes through the reactions catalysed by lysosomal enzymes result in the breakdown of cells and tissues. Many lysosomal enzymes require low pH values for optimal activity. Some of these enzymes are known as acid protease enzymes.

Lysosomes located in cells known as macrophages, and in some other cells, are involved in antigen processing and antigen presentation. Antigen processing and presentation leads to cell to cell interactions within the immune system which triggers release of a set of chemicals called cytokines. Cytokine concentrations in the body are correlated often to many of the clinical signs of inflammation and disease. These clinical signs include heat, swelling, pain, fatigue and lethargy.

An experiment was conducted to assess if fatigue and lethargy could be decreased and motor activity improved by improving the buffering capacity of the cytoplasmic bicarbonate buffer in a group of people diagnosed and suffering with chronic disease.

Improving the buffering capacity of the cytoplasmic bicarbonic buffer would decrease the hydration of carbon dioxide *per se*, would decrease the production of protons from reactions catalysed by carbonic anhydrase enzymes, decrease the pumping of protons by V-type proton pumps, decrease the activities of acid protease enzymes, decrease the activities of lysosomes, decrease antigen processing and presentation, and increase the production of ATP. Fatigue and lethargy would decrease and motor activity would improve. Some of the clinical signs of chronic disease would be alleviated.

A group of twenty three people were chosen who had been diagnosed with having chronic disease. Each person had been suffering from chronic disease for between 3 and 8 years.

The diseases consisted of chronic viral diseases, chronic inflammatory diseases and autoimmune diseases and included rheumatoid arthritis and dermatitis. All people had a history of fatigue and lethargy. In the context of this experiment, fatigue and lethargy were determined as subjective feelings of general exhaustion which were manifested by moderate to severe lack of function. The people consumed aqueous metal bicarbonate solution with a pH value between pH 8.1 and 8.5. The aqueous metal bicarbonate solution contained approximately Mg^{2+} 120 mg per litre, Na^+ 135 mg per litre, K^+ 100 mg per litre and HCO_3^- 1,100 mg per litre. The major component of the solution was magnesium bicarbonate $\text{Mg}(\text{HCO}_3)_2$ 720 mg per litre approximately. Potassium bicarbonate 250 mg per litre was a component of the aqueous metal bicarbonate solution to improve the co-transport of bicarbonate anions into body cells. Consumption of the aqueous metal bicarbonate solution was commenced at half a litre per day and increased by increments over a period of one month to between 2 to 3 litres per day. Consumption occurred on an empty stomach to avoid the loss of bicarbonate by stomach acid (HCl). Consumption occurred in small amounts (300 ml) at set times each day to avoid rapid increases in bicarbonate concentrations of body fluids and to avoid over hydration.


The results of the experiment were delayed but unequivocal. Within 3 to 9 months, all participants in the experiment demonstrated substantial decreases in fatigue and lethargy. All participants demonstrated an increased capacity for mild physical activity; an improvement in motor activity. Function was improved. Those participants with chronic rheumatoid disease (rheumatoid arthritis) demonstrated some decreases in inflammation and some decreases in pain. Those participants with chronic skin disease (dermatitis) demonstrated decreases in inflammation. Those participants with tissue calcification demonstrated decreases in calcium deposits.

The consumption of aqueous metal bicarbonate solution, principally magnesium bicarbonate solution, decreases fatigue and lethargy and improves motor activity in people suffering with chronic disease.

The consumption of aqueous metal bicarbonate solution, principally magnesium bicarbonate solution, decreases clinical signs of inflammation and pain and calcification in people suffering with chronic disease.

An experiment to prevent and to treat the clinical signs of diseases caused by viruses that require proton concentrations for infectivity

Many viruses become infective by utilising high intracellular proton concentrations in host cells. Proton concentrations involved in the infectivity of viruses, and the pathogenesis of 5 viral diseases, derive from the hydration of carbon dioxide catalysed by intracellular carbonic anhydrase enzymes. The production of protons by carbonic anhydrase enzymes can be represented by the equation



10 Protons formed by carbonic anhydrase enzymes are concentrated by intracellular V-type proton pumps and stored in the endosomes and lysosomes of body cells. Functional endosomes and lysosomes maintain internal concentrations of protons which give them internal pH values between pH 3.0 and pH 6.0.

Virus infectivity often requires the activities of acid protease enzymes. Acid protease 15 enzymes are enzymes that function optimally at acidic pH levels. Acid protease enzymes are located in endosomes and lysosomes of body cells.

Influenza viruses require acid protease enzyme activities for their replication and infectivity. The acid proteases of lysosomes and endosomes in body cells act to liberate the nucleic acid (RNA) of the influenza virus from the outer viral membrane.

20 Many viruses contain their own acid-dependent enzymes which utilise proton concentrations in host cells. For example, the acid proteases of lentiviruses are required for virus protein assembly and viral infectivity.

An experiment was instigated with the aim of preventing and treating the clinical signs of diseases caused by viruses that require proton concentrations for infectivity. Natural 25 infections with influenza viruses and flu-like respiratory viruses were taken as the model infections. Influenza is an acute febrile infectious respiratory disease manifested by inflammation of the bronchial mucosa. It is complicated often by bacterial pneumonia. Clinical signs of influenza include initially fever, malaise, headache and muscle pain followed by coughing, sneezing and respiratory tract effusions. Flu-like respiratory 30 viruses cause respiratory diseases manifested generally by clinical signs of less intensity than influenza.

Twenty people were chosen and divided into two equal groups. The control group did not consume aqueous metal bicarbonate solution. The treatment group consumed aqueous metal bicarbonate solution with a pH value between 8.1 and 8.5. The aqueous metal bicarbonate solution contained approximately Mg^{2+} 120 mg per litre, Na^+ 135 mg per 5 litre and HCO_3^- 950 mg per litre. The major component of the solution was magnesium bicarbonate $\text{Mg}(\text{HCO}_3)_2$ 720 mg per litre approximately. Consumption of the aqueous metal bicarbonate solution was commenced at half a litre per day and increased by increments over a period of one month to between 2 to 3 litres per day. Consumption occurred on an empty stomach to avoid the loss of bicarbonate by stomach acid (HCl).

10 Consumption occurred in small amounts (300 ml) at set times each day to avoid rapid increases in bicarbonate concentrations of body fluids and to avoid over hydration.

Aqueous metal bicarbonate solution was consumed by people in the treatment group for 2 years.

People in both groups worked either in child care centres or in homes for the elderly and

15 were exposed to influenza and other respiratory infections over a 2 year period. Clinical signs of influenza and flu-like virus infections were observed and recorded over the 2 years. Results are given below:

Record of influenza and flu-like virus infections over a 2 year period

	Control Group	Treatment Group
Influenza		
•Number of infections	8	2
•Duration of symptoms	5 to 10 days	2 to 3 days
•Severity of symptoms (0 to 4)	4	1 to 2
Flu-like viruses		
•Number of infections	15	3
•Duration of symptoms	3 to 7 days	2 to 3 days
•Severity of symptoms (0 to 4)	2 to 4	1 to 2

People consuming aqueous metal bicarbonate solution had a lower prevalence of the

20 clinical signs of influenza and flu-like virus infections than people not consuming aqueous metal bicarbonate solution. People consuming aqueous metal bicarbonate solution had

less severe symptoms and shorter duration of symptoms than people not consuming aqueous metal bicarbonate solution.

The consumption of aqueous metal bicarbonate solution, principally magnesium bicarbonate solution, decreases the prevalence of the clinical signs of diseases caused by viruses that require proton concentrations for infectivity.

The consumption of aqueous metal bicarbonate solution, principally magnesium bicarbonate solution, decreases the severity and the duration of the clinical signs of diseases caused by viruses that require proton concentrations for infectivity.

Example 8

10 Suitable formulation and range of parameters and range of administration volumes for the aqueous metal bicarbonate solution

A suitable formulation for the aqueous metal bicarbonate solution contains $\text{Mg}(\text{HCO}_3)_2$ 720 mg per litre approximately (Mg^{2+} 120 mg per litre and HCO_3^- 600 mg per litre approximately). The formulation usually contains also NaHCO_3 485 mg per litre approximately (Na^+ 135 mg per litre and HCO_3^- 350 mg per litre approximately). The pH of the formulation is pH 8.3. The formulation is stored either at 5°C to 10°C at 1 atmosphere in a sealed container or at higher temperatures at higher pressures. The formulation is administered or consumed by a mammalian (typically human) user in 300 ml doses approximately 6 to 10 times per day usually on an empty stomach at approximately equal time intervals throughout the day. Usually the total amount of formulation usually administered is 1.8 to 3 litres per day.

The parameters of a suitable formulation for the aqueous metal bicarbonate solution may be conveniently represented as follows:

Typical parameter range	Specific parameters
$\text{Mg}(\text{HCO}_3)_2$ 150 mg/litre to saturation solubility per litre	$\text{Mg}(\text{HCO}_3)_2$ 720 mg/litre (Mg^{2+} 120 mg/litre and HCO_3^- 600 mg/litre)
NaHCO_3 0 to 1,000 mg/litre	NaHCO_3 485 mg/litre (Na^+ 135 mg/litre and HCO_3^- 350 mg/litre)
pH 8.0 to 8.5	pH 8.3

0°C to 10°C at 1 atmosphere	5°C to 10°C at 1 atmosphere
300 ml dose approximately 1 to 15 times per day	300 ml dose approximately 6 to 10 times per day

There exists a range of combinations and concentrations of metal cations that may be included in the suitable formulation of the aqueous metal bicarbonate solution. There exists a range of anions (other than bicarbonate) that may be included in stoichiometric amounts with metal cations in the suitable formulation of the aqueous metal bicarbonate solution. There exists a range of concentrations of bicarbonate anions that may be included in stoichiometric amounts with metal cations in the suitable formulation of the aqueous metal bicarbonate solution.

A typical range of parameters that may be combined and included in the suitable formulation of the aqueous metal bicarbonate solution may be conveniently represented as follows:

Final Concentration in Solution

	Typical parameter range	Specific parameters
Mg ²⁺	50 to 500 mg/litre	120 to 300 mg/litre
Na ⁺	50 to 1,000 mg/litre	120 to 300 mg/litre
K ⁺	50 to 500 mg/litre	120 to 300 mg/litre
Ca ²⁺	50 to 500 mg/litre	120 to 300 mg/litre
HCO ₃ ⁻	200 to 3,000 mg/litre	600 to 2,000 mg/litre
Anions other than HCO ₃ ⁻ (eg. Cl ⁻ , SO ₄ ²⁻)	Stoichiometric with metal cation concentrations	

There exists a range of pH values for the suitable formulation of aqueous metal bicarbonate solution (that includes cation and anion parameter ranges above). The pH values may be conveniently represented as follows:

Typical pH range	Specific pH range
pH 7.5 to 8.8	pH 8.0 to 8.5

15 Above pH 8.5, the solution tends to become cloudy due to the formulation of metal carbonates (CO₃²⁻).

There exists a range of physical parameters for the suitable formulation of aqueous metal bicarbonate solution (that includes cation and anion parameter ranges above). The physical parameters may be conveniently represented as follows:

Typical parameter range	Specific parameter range
0°C to 30°C	5°C to 10°C
1 to 3 atmospheres	1 atmosphere

Above 10°C, carbon dioxide leaves solution and the solution tends to become cloudy (with 5 time) due to the formation of metal carbonates (CO_3^{2-}). This may be controlled by increasing the pressure on the solution.

There exists a range of volumes of administration for the suitable formulation of metal bicarbonate solution (that includes cation and anion parameter ranges above). The volume of aqueous metal bicarbonate solution administered depends on the purpose for the 10 administration.

The administered volumes of aqueous metal bicarbonate solution may be conveniently represented as follows:

Purpose for administration of aqueous metal bicarbonate solution	Typical volume range administered to 70 kg human
Increased longevity and delay in senescence	1 to 2 litres per day (normal physiological volume requirement for water intake)
Prevention of degenerative diseases	1 to 2 litres per day (normal physiological volume requirement for water intake)
Treatment of osteoarthritis	1.8 to 3.0 litres per day
Treatment of chronic disease	1.8 to 3.0 litres per day
Maintain and improve motor activity	1.8 to 3.0 litres per day
Decrease fatigue and lethargy	1.8 to 3.0 litres per day
Prevention and treatment of influenza and other acid-dependent viral diseases	1.8 to 3.0 litres per day

Aqueous metal bicarbonate solution is administered typically in 300 ml doses approximately 6 to 10 times per day at equal time intervals throughout the day.

Typically, the solution is allowed to stand prior to consumption until the solution reaches 15°C to 20°C (cool room temperature).

- 5 A suitable formulation for the aqueous metal bicarbonate solution contains $\text{Mg}(\text{HCO}_3)_2$ 720 mg per litre at pH 8.3. The formulation usually contains also NaHCO_3 485 mg per litre. The aqueous metal bicarbonate solution may contain a range of cations and anions within a pH range pH 7.5 to 8.8. The aqueous metal bicarbonate solution is administered in volumes ranging from 1 to 3 litres per day.

10

Example 9

- Osteoarthritis is a slowly progressive degeneration of the joints of the hands and large weight-bearing joints (hips and knees). Osteoarthritis is common in post menopausal women. Osteoarthritis is characterised by pain, enlargement of joints and limitation of joint movements. The linings of osteoarthritic joints show a moderate to marked degree of inflammation. The principle pathological changes associated with osteoarthritis are destruction of joint cartilage and neoformations of bone at joint margins (osteophytes). In osteoarthritis, destruction of joint cartilage is caused by acid protease enzymes (and other enzymes) derived often from the lysosomes of cartilage cells (chondrocytes), inflammatory cells and other cells.
- 20 For an experimental trial, a group of post menopausal women were chosen who had clinical signs of osteoarthritis in the joints of their hands. The osteoarthritic joints included the distal and proximal interphalangeal joints of the fingers and the carpometacarpal joint of the thumbs. In all cases, loss of joint function was moderate to severe.
- 25 In all cases, the women suffered pain, swelling of the fingers and loss of joint movement. Mucous cysts were associated with distal joint osteoarthritis. Lateral deformities occurred in some proximal joints with severe loss of joint function. Women with affected thumbs had considerable loss of function and considerable pain. Many hands were "claw-like" in appearance. Women consumed magnesium bicarbonate solution, with added sodium bicarbonate. The women consumed between 2 to 3 litres of bicarbonate solution per day.
- 30 In this solution, the magnesium concentration was approximately 120mg per litre, sodium concentration was approximately 135mg per litre and bicarbonate concentration was

approximately 950mg per litre. Consumption was commenced at half a litre per day and increased by increments over a period of one month to 2 to 3 litres per day. Consumption occurred on an empty stomach to avoid the loss of bicarbonate by stomach acid. Consumption occurred in small amounts (300mL) at set times each day to avoid rapid
5 increases in bicarbonate concentrations of body fluids.

In all cases, there were remissions in the clinical signs of osteoarthritis which were quantifiable by standard tests of movement, flexibility and strength. The participants showed considerable increases in joint functions and decreases in acute and chronic joint swellings. The "stabbing" pain of osteoarthritis was alleviated. Some participants had
10 remissions of inflammation and arthritis to the stage where many chronic swellings were no longer observable and joint mobilities and functions were restored. However, these improvements were maintained only with the continued consumption of bicarbonate anions. Once the consumption of bicarbonate anions ceased, clinical signs of inflammation began to reappear often within a week.

15 The participants commented on an absence of lethargy and the presence of a feeling of well-being. Magnesium bicarbonate alleviated the pain and swelling associated with osteoarthritis.

None of the participants demonstrated any clinical signs of influenza or other respiratory viral infections over the two year period of the trial. This occurred despite several of the
20 participants working in situations where exposure to viral infections was high (nursing homes and child care centres).

Example 10

Influenza is an acute febrile infectious respiratory disease manifested by inflammation of the bronchial mucosa. Influenza is complicated often by bacterial pneumonia which may
25 be fatal.

For an experimental trial, a group of men and women were chosen who worked in situations where exposure to the influenza virus was likely to occur (nursing homes and child care centres). Each person in the experimental group consumed magnesium bicarbonate solution, with added sodium bicarbonate. Each person consumed between 2
30 to 3 litres of bicarbonate solution per day. In this solution, the magnesium concentration was approximately 120mg per litre, sodium concentration was approximately 135mg per litre and bicarbonate concentration was approximately 950mg per litre.

Consumption was commenced at half a litre per day and increased by increments over a period of one month to 2 to 3 litres per day. Consumption occurred on an empty stomach to avoid the loss of bicarbonate by stomach acid. Consumption occurred in small amounts (300mL) at set times each day to avoid rapid increases in bicarbonate concentrations of 5 body fluids.

Over the two year period of the experimental trial, no person in the experimental group showed any clinical signs of influenza.

Example 11

A suitable formulation for the aqueous metal bicarbonate solution contains 720mg 10 $\text{Mg}(\text{HCO}_3)_2$ per litre (120mg Mg^{2+} per litre and 600mg HCO_3^- per litre approximately) and 485mg NaHCO_3 per litre (135mg Na^+ per litre and 350mg HCO_3^- per litre approximately). The pH of this formulation is pH 8.3. This formulation is stored at 5 to 10°C at 1 atmosphere or it can be stored at higher temperatures at higher pressures. This formulation is administered in 300mL doses approximately 6 to 10 times per day. That is 15 the amount of formulation usually administered per day is 1.8 to 3 litres per day.

The parameters of the formulation may be conveniently represented as follows:

Typical Parameter Range	Specific Parameters
100mg - saturation solubility $\text{Mg}(\text{HCO}_3)_2$ per litre	720mg $\text{Mg}(\text{HCO}_3)_2$ per litre (120mg Mg^{2+} and 600mg HCO_3^- per litre)
0-1000mg NaHCO_3 per litre	485mg NaHCO_3 per litre (135mg Na^+ and 350mg HCO_3^- per litre)
pH 8-8.6, typically 8-8.5	pH 8.3
0-10°C @ 1 atmosphere	5-10°C @ 1 atmosphere
300mL dose approximately 1 to 20 times per day	300mL dose approximately 6 to 10 times per day

Example 12

Mitochondria are inefficient if they cannot maintain the necessary production of ATP for maintenance of essential cell processes and cell functions. This inefficiency is due often 20 to functional damage to the mitochondrial inner membrane and other mitochondrial molecules. Inefficient mitochondria are not able to maintain normal carbon, electron and proton fluxes.

However, in middle age, carbon and electron fluxes may be maintained by the synthesis of fatty acids in the cytoplasm of body cells. In body cells of the middle aged, fatty acids can be regarded as carbon and electron sinks necessary for the maintenance of essential fluxes; that is, for the maintenance of essential life processes. The synthesis of fatty acids

5 utilises ATP. However, the fluxes for production of ATP in mitochondria are decreased in middle age. There is a consequent 'energy' deficit. The middle aged are flat and lethargic relative to the young, though they can be regarded as fat by necessity; the necessity of staying alive. In addition to utilisation of cell ATP, the synthesis of fatty acids in body cells adds to the carbon dioxide load of the cells and adds to concentrations of

10 intracellular protons. This occurs because the series of chemical reactions that synthesise fatty acids results in a net utilisation of bicarbonate anions and a net production of carbon dioxide and protons. For example, each molecule of the fatty acid palmitate that is synthesised by cells utilises seven molecules of ATP and seven bicarbonate anions and produces seven molecules of carbon dioxide and seven protons. Of course, fatty acids are

15 oxidised continuously from fat stores in the body which produces even more carbon dioxide. When excess calories are consumed at any chronological age (and converted to fatty acids) the overall carbon dioxide load is considerable. Caloric restricted rodents avoid this extra carbon dioxide load and, as a consequence, they live longer lives with delays in the onset of degenerative diseases.

20 A trial involving people consuming bicarbonate anions in water was conducted. These people were middle aged and overweight and complained of tiredness and lethargy. No control group was maintained for the duration of the trial (people in an initial control group were unable to consume the volumes of soft water required). People involved in the trial were given a series of lectures on the biochemistry of mitochondrial processes. They

25 were requested to decrease their food (calorie) intake considerably and to avoid dietary fats. Excessive aerobic exercise was not recommended due to the consequent increase in hunger it produces, the large increase in carbon dioxide concentrations that occur with increased aerobic muscle activity and the damage excess activity does to inefficient mitochondria. Indeed, active muscle cells contain mitochondria with most nucleic acid

30 damage relative to other body cells.

The trial consisted of each person consuming between two to three litres per day of a mixture of magnesium bicarbonate and sodium bicarbonate in water. Bicarbonate concentration was established at a maximum of one gram per litre. (This concentration of

bicarbonate is well within the concentrations in several water sources utilised for human consumption in Europe. In these waters however, the bicarbonate is in the form of calcium and sodium bicarbonate and pH values often are not very alkaline.) Consumption was commenced at half a litre per day and increased by increments over a period of one 5 month to the maximum consumption. This start-up schedule avoided any gastrointestinal side effects due to the smooth muscle relaxation properties of magnesium. Capillary dilation in the face was apparent in several people (which was interpreted by those affected as rosy, healthy cheeks). The capillary dilation may have been due to magnesium or may have been due to a decrease in activity of renin which is an acid protease enzyme 10 from the kidney that is central to the control of blood pressure. (Renin exerts its effects ultimately by constriction of small blood vessels.)

Each participant in the trial was advised to consume the bicarbonate solution 'on an empty stomach'. Consumption in this manner avoided the mixing of bicarbonate anions with stomach acid which would have resulted in the loss of bicarbonate. Advice was given also 15 to consume the solution in small amounts at set times each day. Consumption in this manner avoided a rapid increase in the bicarbonate concentrations of body fluids.

The results were unequivocal. Body weight was lost at about one half of a kilogram per week after the initial start-up period was completed. Other beneficial effects (more important than weight loss to the participants in the trial) included the absence of lethargy, 20 the presence of a feeling of well-being (mild euphoria) and the increased capacity for mild physical activity. The participants all commented that their 'energy levels' had improved and that their outlook on life had consequently become more positive.

Example 13

A heart muscle cell contains mitochondria that occupy one quarter of the cell volume. It is 25 natural to expect the heart to be rich in mitochondria when one considers the workload of the heart and its subsequent requirement for 'energy'. The consumption of magnesium bicarbonate may assist in maintaining efficient mitochondria in heart muscle cells. In the presence of bicarbonate anions, mitochondrial efficiency in heart muscle cells is maintained by processes which include decreases in proton leaks across inner mitochondrial 30 membranes, establishment of proton circuits independent of proton leaks and maintenance of alkaline pH values in mitochondrial matrixes. In the presence of bicarbonate anions, mitochondrial damage and mitochondrial failure are decreased. Efficient mitochondria in

heart muscle cells maintain ATP production so that the heart remains functional as a vital organ.

There are further beneficial effects to the cardiovascular system resulting from the consumption of magnesium bicarbonate. First, decreased fatty acid synthesis in body cells results in lowered body weight with subsequent reduction in high blood pressure to a normal value. Indeed, these effects were observed in people participating in the trial reported in Example 5. Second, lysosomal enzyme damage to ischaemic heart muscle may be prevented or decreased.

There is unequivocal evidence of correlation that heart disease, senescence, degenerative diseases and death are caused by inefficient mitochondria. Though inefficient mitochondria may not be the only cause of these conditions. There is sufficient evidence however that senescence and general degeneration in all species studied, from fungi to humans, are correlated to the damage to the complex molecules of the mitochondria. For example, the mitochondria of aged people carry nucleic acid and protein defects not observed in the mitochondria of young people. This is true particularly of the mitochondria in muscle, heart and brain cells. Accordingly, it has been proposed that several chronic diseases common in old age may be related to mitochondrial failure, including heart disease, late-onset diabetes and Parkinson's and Alzheimer's diseases. That is, the gradual loss of efficiency of cell mitochondria results in a diminution of the functional capacity of body cells with pathological consequences.

A progressive decline in organ function is characteristic of old age. Some of the changes that occur as people grow older are:

1. The ability to focus the eyes decreases and response time to stimuli becomes slower.
2. Cancers in epithelial tissues (skin, lung, colon, mammary gland) become more common.
3. Heart disease and widely disseminated atherosclerosis occur.
4. Osteoporosis and bone and joint pathology occur.
5. Thermoregulation becomes impaired.
6. There is a decline in the ability of body organs such as the reproductive organs, lungs, glandular tissues and kidneys to maintain their specialised life processes.
7. There is a reduced capacity for surviving haemorrhage.

8. There is an increase in autoimmune diseases and chronic inflammatory diseases.

Relative to their life span of around three years, mice develop heart disease, kidney disease, arthritis and cancer at similar stages in life to humans. In other words, both senescence and the similar degenerative diseases of mice and humans are correlated to the 5 chronological age of each species but cannot be linked causally to chronological age *per se*. It has been observed in many experiments that rodents fed caloric restricted diets suffer less from the above diseases than control animals. It has been observed also that careful necropsies on diet-restricted rodents often do not reveal any gross or microscopic pathology.

10 Mitochondrial proton leak is a contributing cause of mitochondrial inefficiency and hence a source of senescence and degenerative diseases in short-lived rodents. Mitochondrial proton leak may be a source of senescence and degenerative diseases in humans. Because the proton leak in humans is only a fraction of the leak in rodents, humans develop senescence and degenerative diseases at a later chronological age than rodents.

15 It may be possible to prevent inefficient processes in mitochondria in order to extend longevity and to delay degenerative diseases. Exogenous sources of appropriate bicarbonate anions can be absorbed by body cells to maintain intracellular alkaline pH values and that alkaline pH values maintain mitochondrial respiration rates and maintain effective proton concentration gradients across inner mitochondrial membranes.

20 Intracellular alkaline pH values appear necessary for optimum activities of many enzymes in body cells. These enzymes include polymerases, phosphofructokinase and carbonic anhydrase. Enzymes permit life processes to be perpetuated. Therefore, maintenance of enzyme activities by maintenance of intracellular alkaline pH values may assist in the perpetuation of longevity.

25 The long-term provision of appropriate bicarbonate anions to body cells maintains efficient mitochondrial function, maintains the DNA polymerase activity of mitochondria and therefore maintains the integrity of mitochondrial DNA. This latter effect results in accurate syntheses of the complex functional molecules that are involved in electron fluxes in mitochondria. Of course, bicarbonate anions will decrease the proton load *per se* in 30 body cells with a subsequent decrease in proton damage to other cell molecules and a decrease in the tendency of oxidation reactions to occur. Decreased oxidations result in

decreased DNA mutations and decreased amino acid oxidations. Degenerative diseases are delayed.

Industrial Applicability

An aqueous metal bicarbonate solution of the invention can be readily utilised in medicine
5 to prevent and to treat certain inflammatory diseases, degenerative diseases and viral diseases in mammals.

The claims defining the invention are as follows:

1. An aqueous neutral to mildly alkaline metal bicarbonate solution, comprising magnesium bicarbonate dissolved in the solution, said magnesium bicarbonate comprising bicarbonate anions and magnesium cations, and a pH adjusting agent in the solution in an amount whereby the solution is at a neutral to mildly alkaline pH.
2. The aqueous neutral to mildly alkaline metal bicarbonate solution of claim 1, wherein the bicarbonate anions are 150mg to 3500mg per litre of the solution.
3. The aqueous neutral to mildly alkaline metal bicarbonate solution of claim 1 or 2, wherein the magnesium cations are 30mg to 500mg per litre of the solution.
4. The aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 3, wherein a stoichiometric concentration of magnesium cations is in association with the bicarbonate anions.
5. The aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 4, wherein the pH adjusting agent is selected from the group consisting of carbon dioxide, hydrated carbon dioxide, carbonic acid and any mixture thereof.
6. The aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 5, wherein the aqueous metal bicarbonate solution has a pH 7 to 8.6.
7. The aqueous metal bicarbonate solution of any one of claims 1 to 5, wherein the aqueous neutral to mildly alkaline metal bicarbonate solution has a pH of 8.0 to 8.6.
8. The aqueous metal bicarbonate solution of any one of claims 1 to 7, wherein the aqueous neutral to mildly alkaline metal bicarbonate solution has a temperature in the range selected from the group consisting of 0 to 25°C, 0 to 20°C, 0.5 to 25°C, 0.5 to 20°C, 0.5 to 15°C, 0.5 to 10°C, 0.5 to 9°C, 0.5 to 8°C, 0.5 to 7°C, 1 to 20°C, 1 to 15°C, 1 to 10°C, 1.5 to 20°C, 1.5 to 15°C, 1.5 to 10°C, 2 to 20°C, 2 to 15°C, 2 to 10°C, 3 to 20°C, 3 to 15°C, 4 to 20°C, 4 to 15°C, 4 to 10°C, 5 to 20°C, 5 to 15°C, 6 to 20°C, 6 to 15°C, 6 to 10°C, 7 to 20°C, 7 to 15°C, 7 to 10°C, 8 to 20°C, 8 to 15°C, 8 to 10°C, 9 to 20°C, 9 to 15°C, 9 to 10°C, 10 to 15°C, 0 to 15°C, 0 to 10°C, 3°C to 10°C and 5°C to 10°C.
9. An aqueous neutral to mildly alkaline metal bicarbonate solution, comprising magnesium bicarbonate dissolved in the solution, substantially as herein described with reference to any one of Examples 1, 3, 4, 5, 6, 7, 9 and 13.
10. A solution for preventing certain inflammatory diseases in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 wherein the magnesium bicarbonate is in an amount effective to prevent said diseases.



11. A solution for treating certain inflammatory diseases in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 wherein the magnesium bicarbonate is in an amount effective to treat said diseases.

5 12. A solution for preventing degenerative diseases in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 wherein the magnesium bicarbonate is in an amount effective to prevent said diseases.

13. A solution for treating degenerative diseases in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9
10 wherein the magnesium bicarbonate is in an amount effective to treat said diseases.

14. A solution for preventing certain viral diseases in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 wherein the magnesium bicarbonate is in an amount effective to prevent said diseases.

15 15. A solution for treating certain viral diseases in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 wherein the magnesium bicarbonate is in an amount effective to treat said diseases.

16. A solution for decreasing senescence in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 wherein the magnesium bicarbonate is in an amount effective to decrease senescence.

20 17. A solution for treating senescence in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 wherein the magnesium bicarbonate is in an amount effective to treat senescence.

18. A solution for increasing longevity in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 wherein
25 the magnesium bicarbonate is in an amount effective to increase longevity.

19. A solution for scavenging protons in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 wherein the magnesium bicarbonate is in an amount effective to scavenge protons.

30 20. A solution for decreasing proton concentrations in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 wherein the magnesium bicarbonate is in an amount effective to decrease proton concentrations.

21. A solution for decreasing inflammation in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9, wherein the magnesium bicarbonate is in an amount effective to decrease inflammation.



22. A solution for decreasing inflammatory conditions in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9, wherein the magnesium bicarbonate is in an amount effective to decrease inflammatory conditions.

5 23. A solution for increasing motor activity in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 wherein the magnesium bicarbonate is in an amount effective to increase motor activity.

24. A solution for decreasing fatigue in a mammal, comprising the aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 wherein
10 the magnesium bicarbonate is in an amount effective to decrease fatigue.

25. A process of preparing an aqueous neutral to mildly alkaline metal bicarbonate solution comprising bicarbonate anions and metal cations, which process comprises reacting a compound selected from the group consisting of metal carbonate, metal carbonate hydroxide, metal oxide, metal hydroxide and any mixture thereof with an
15 effective concentration of a pH adjusting agent to produce the aqueous neutral to mildly alkaline metal bicarbonate solution, wherein the pH adjusting agent is present in an amount whereby the solution is at a neutral to mildly alkaline pH.

26. The process of preparing an aqueous neutral to mildly alkaline metal bicarbonate solution of claim 25, wherein the pH adjusting agent is selected from the
20 group consisting of carbon dioxide, hydrated carbon dioxide, carbonic acid and any mixture thereof.

27. The process of preparing an aqueous neutral to mildly alkaline metal bicarbonate solution of claim 25 or 26, wherein the metal bicarbonate is selected from the group consisting of magnesium bicarbonate, sodium bicarbonate, a mixture of sodium
25 bicarbonate and magnesium bicarbonate, potassium bicarbonate, a mixture of potassium bicarbonate and magnesium bicarbonate, calcium bicarbonate, a mixture of calcium bicarbonate and magnesium bicarbonate, lithium bicarbonate, and a mixture of lithium bicarbonate and magnesium bicarbonate.

28. The process of preparing an aqueous neutral to mildly alkaline metal
30 bicarbonate solution of any one of claims 25 to 27, wherein the bicarbonate anions are 150mg to 3500mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution.

29. The process of preparing an aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 25 to 28, wherein the metal cations are 30mg to 500mg per litre of aqueous neutral to mildly alkaline metal bicarbonate solution



30. The process of preparing an aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 25 to 29, wherein the aqueous neutral to mildly alkaline metal bicarbonate solution has a pH of 7 to 8.6.

31. The process of preparing an aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 25 to 30, wherein the aqueous metal bicarbonate solution has a pH of 8.0 to 8.6.

32. The process of an aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 25 to 30, wherein the aqueous metal bicarbonate solution has a temperature in the range selected from the group consisting of 0 to 25°C, 0 to 20°C, 0.5 to 25°C, 0.5 to 20°C, 0.5 to 15°C, 0.5 to 10°C, 0.5 to 9°C, 0.5 to 8°C, 0.5 to 7°C, 1 to 20°C, 1 to 15°C, 1 to 10°C, 1.5 to 20°C, 1.5 to 15°C, 1.5 to 10°C, 2 to 20°C, 2 to 15°C, 2 to 10°C, 3 to 20°C, 3 to 15°C, 4 to 20°C, 4 to 15°C, 4 to 10°C, 5 to 20°C, 5 to 15°C, 6 to 20°C, 6 to 15°C, 6 to 10°C, 7 to 20°C, 7 to 15°C, 7 to 10°C, 8 to 20°C, 8 to 15°C, 8 to 10°C, 9 to 20°C, 9 to 15°C, 9 to 10°C, 10 to 15°C, 0 to 15°C, 0 to 10°C, 3°C to 10°C and 5°C to 10°C.

33. A process of preparing an aqueous neutral to mildly alkaline metal bicarbonate solution, which process is substantially as herein described with reference to any one of the Examples.

34. An aqueous neutral to mildly alkaline metal bicarbonate solution whenever prepared by the process of any one of claims 25 to 33.

35. A combination comprising a substantially stable aqueous neutral to mildly alkaline metal bicarbonate solution, comprising magnesium bicarbonate dissolved in the solution, said magnesium bicarbonate comprising bicarbonate anions and magnesium cations, and a pH adjusting agent in the solution in an amount whereby the solution is at a neutral to mildly alkaline pH, in combination with a stabilising agent in an amount effective to maintain and stabilise the bicarbonate anions in the neutral to mildly alkaline solution.

36. A combination comprising a substantially stable aqueous neutral to mildly alkaline metal bicarbonate solution, substantially as herein described with reference to any one of Examples 1, 3, 4, 5, 6, 7, 9 or 13.

37. A method of preventing certain inflammatory diseases in a mammal in need of such prevention comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 10 or a combination of claim 35 or 36.



38. A method of treating certain inflammatory diseases in a mammal in need of such treatment comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 11 or a combination of claim 35 or 36.

39. A method of preventing degenerative diseases in a mammal in need of such prevention comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 12 or a combination of claim 35 or 36.

40. A method of treating degenerative diseases in a mammal in need of such treatment comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 13 or a combination of claim 35 or 36.

41. A method of preventing certain viral diseases in a mammal in need of such prevention comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 14 or a combination of claim 35 or 36.

42. A method of treating certain viral diseases in a mammal in need of such treatment comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 15 or a combination of claim 35 or 36.

43. A method of decreasing senescence in a mammal comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 16 or a combination of claim 35 or 36.

44. A method of treating senescence in a mammal comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 17 or a combination of claim 35 or 36.

45. A method of increasing longevity in a mammal comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 18 or a combination of claim 35 or 36.

46. A method of scavenging protons in a mammal comprising administering to said mammal an effective amount of the aqueous neutral to mildly alkaline metal



bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 19 or a combination of claim 35 or 36.

47. A method of decreasing proton concentrations in a mammal by altering carbonic anhydrase enzyme reactions in said mammal comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 20 or a combination of claim 35 or 36.

48. A method of decreasing inflammation in a mammal comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 21 or a combination of claim 35 or 36.

49. A method of decreasing inflammatory conditions in a mammal comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 22 or a combination of claim 35 or 36.

50. A method of increasing motor activity in a mammal comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 23 or a combination of claim 35 or 36.

51. A method of decreasing fatigue in a mammal comprising administering to said mammal an effective amount of an aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 24 or a combination of claim 35 or 36.

52. The method of any one of claims 37 to 51, wherein said mammal is human and said aqueous neutral to mildly alkaline metal bicarbonate solution is administered to said human on an empty stomach.

53. Use of bicarbonate, magnesium cations and a pH adjusting agent of any one of claims 1 to 9 or 34 or a combination of claim 35 or 36 for the manufacture of a medicament to prevent certain inflammatory diseases in a mammal in need of such prevention.

54. Use of bicarbonate, magnesium cations and a pH adjusting agent of any one of claims 1 to 9 or 34 or a combination of claim 35 or 36 for the manufacture of a medicament to treat certain inflammatory diseases in a mammal in need of such treatment.



55. Use of bicarbonate, magnesium cations and a pH adjusting agent of any one of claims 1 to 9 or 34 or a combination of claim 35 or 36 for the manufacture of a medicament to prevent degenerative diseases in a mammal in need of such prevention.

56. Use of bicarbonate, magnesium cations and a pH adjusting agent of any one of claims 1 to 9 or 34 or a combination of claim 35 or 36 for the manufacture of a medicament to treat degenerative diseases in a mammal in need of such treatment.

57. Use of bicarbonate, magnesium cations and a pH adjusting agent of any one of claims 1 to 9 or 34 or a combination of claim 35 or 36 for the manufacture of a medicament to prevent certain viral diseases in a mammal in need of such prevention.

58. Use of bicarbonate, magnesium cations and a pH adjusting agent of any one of claims 1 to 9 or 34 or a combination of claim 35 or 36 for the manufacture of a medicament to treat certain viral diseases in a mammal in need of such treatment.

59. Use of bicarbonate, magnesium cations and a pH adjusting agent of any one of claims 1 to 9 or 34 or a combination of claim 35 or 36 for the manufacture of a medicament to decrease senescence in a mammal.

60. Use of bicarbonate, magnesium cations and a pH adjusting agent of any one of claims 1 to 9 or 34 or a combination of claim 35 or 36 for the manufacture of a medicament to treat senescence in a mammal.

61. Use of bicarbonate, magnesium cations and a pH adjusting agent of any one of claims 1 to 9 or 34 or a combination of claim 35 or 36 for the manufacture of a medicament to increase longevity in a mammal.

62. Use of bicarbonate, magnesium cations and a pH adjusting agent of any one of claims 1 to 9 or 34 or a combination of claim 35 or 36 for the manufacture of a medicament to scavenge protons in a mammal.

63. Use of bicarbonate, magnesium cations and a pH adjusting agent of any one of claims 1 to 9 or 34 or a combination of claim 35 or 36 for the manufacture of a medicament to decrease proton concentrations in a mammal.

64. Use of bicarbonate, magnesium cations and a pH adjusting agent of any one of claims 1 to 9 or 34 or a combination of claim 35 or 36 for the manufacture of a medicament to decrease inflammation in a mammal.

65. Use of bicarbonate, magnesium cations and a pH adjusting agent of any one of claims 1 to 9 or 34 or a combination of claim 35 or 36 for the manufacture of a medicament to decrease inflammatory conditions in a mammal.



66. Use of bicarbonate, magnesium cations and a pH adjusting agent of any one of claims 1 to 9 or 34 or a combination of claim 35 or 36 for the manufacture of a medicament to increase motor activity in a mammal.

67. Use of bicarbonate, magnesium cations and a pH adjusting agent of any one of claims 1 to 9 or 34 or a combination of claim 35 or 36 for the manufacture of a medicament to decrease fatigue in a mammal.

68. The use of any one of claims 53 to 67, wherein said mammal is human and said bicarbonate, magnesium cation and pH adjusting agent or combination is administered to said human on an empty stomach.

69. An aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 10 or a combination of claim 35 or 36 when used to prevent certain inflammatory diseases in a mammal in need of such prevention.

70. An aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 11 or a combination of claim 35 or 36 when used to treat certain inflammatory diseases in a mammal in need of such treatment.

71. An aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 12 or a combination of claim 35 or 36 when used to prevent degenerative diseases in a mammal in need of such prevention.

72. An aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 13 or a combination of claim 35 or 36 when used to treat degenerative diseases in a mammal in need of such treatment.

73. An aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 14 or a combination of claim 35 or 36 when used to prevent certain viral diseases in a mammal in need of such prevention.

74. An aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 15 or a combination of claim 35 or 36 when used to treat certain viral diseases in a mammal in need of such treatment.

75. An aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 16 or a combination of claim 35 or 36 when used to decrease senescence in a mammal.

76. An aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 17 or a combination of claim 35 or 36 when used to treat senescence in a mammal.



77. An aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 18 or a combination of claim 35 or 36 when used to increase longevity in a mammal.

78. An aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 19 or a combination of claim 35 or 36 when used to scavenge protons in a mammal.

79. An aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 20 or a combination of claim 35 or 36 when used to decrease proton concentrations in a mammal.

80. An aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 21 or a combination of claim 35 or 36 when used to decrease inflammation in a mammal.

81. An aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 22 or a combination of claim 35 or 36 when used to decrease inflammatory conditions in a mammal.

82. An aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 23 or a combination of claim 35 or 36 when used to increase motor activity in a mammal.

83. An aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 1 to 9 or 34 or a solution of claim 24 or a combination of claim 35 or 36 when used to decrease fatigue in a mammal.

84. The aqueous neutral to mildly alkaline metal bicarbonate solution of any one of claims 69 to 83, wherein said mammal is human and said aqueous neutral to mildly alkaline metal bicarbonate solution is administered to said human on an empty stomach.

85. The medicament whenever manufactured by the use of any one of claims 53 to 68.

Dated 11 May 2001

Macquarie Veterinary Supplies Pty Ltd

Patent Attorneys for the Applicant/Nominated Person

SPRUSON & FERGUSON

30



ABSTRACT

An aqueous neutral to mildly alkaline metal bicarbonate solution is disclosed. The solution comprises metal bicarbonate dissolved in the solution, the metal bicarbonate comprising bicarbonate anions and metal cations. In addition there is a pH adjusting agent in the solution in an amount whereby the solution is at a neutral to mildly alkaline pH. Also disclosed is a process of preparing an aqueous neutral to mildly alkaline metal bicarbonate solution comprising bicarbonate anions and metal cations. The process comprises reacting a compound selected from the group consisting of metal carbonate, metal carbonate hydroxide, metal oxide, metal hydroxide and any mixture thereof with an effective concentration of a pH adjusting agent to produce the aqueous neutral to mildly alkaline metal bicarbonate solution, wherein the pH adjusting agent is present in an amount whereby the solution is at a neutral to mildly alkaline pH. Further disclosed are a method of preventing and/or treating certain inflammatory diseases and/or degenerative diseases in a mammal, a method of preventing and/or treating certain viral diseases in a mammal, a method of decreasing and/or treating senescence and/or of increasing longevity in a mammal, a method of scavenging protons in a mammal, a method of decreasing proton concentrations in a mammal by altering carbonic anhydrase enzyme reactions in said mammal, a method of decreasing inflammation and/or inflammatory conditions in a mammal and a method of increasing motor activity and/or decreasing fatigue in a mammal.




FIGURE 1

FIGURE 2




FIGURE 3



U
S
A
S
E
O

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.